

EVOLVING CONCEPTS OF CHONDROGENIC DIFFERENTIATION: HISTORY, STATE-OF-THE-ART AND FUTURE PERSPECTIVES

X. Tang^{1,†}, L. Fan^{1,†}, M. Pei², L. Zeng³ and Z. Ge^{1,4*}

¹ Department of Biomedical Engineering, College of Engineering, Peking University, Beijing, China

² Stem Cell and Tissue Engineering Laboratory, Department of Orthopaedics, West Virginia University, Morgantown, West Virginia, USA

³ Department of Integrative Physiology and Pathobiology, Tufts University School of Medicine, Boston, Massachusetts, USA

⁴ Arthritis Clinic and Research Centre, Peking University People's Hospital, Beijing, China

† These authors contributed equally to this work

Abstract

As a cell source, multipotent mesenchymal stromal cells or mesenchymal stem cells (MSCs) are promising candidates for chondrogenic differentiation and subsequent cartilage regeneration. From previous literature, it is known that chondrogenic differentiation of MSCs inevitably leads to hypertrophy and subsequent endochondral ossification. In this review, we examine the history of currently established protocols of chondrogenic differentiation and elaborate on the roles of individual components of chondrogenic differentiation medium. We also summarise the effects of physical, chemical and biological factors involved, and propose potential strategies to differentiate MSCs into articular chondrocytes with homogenous mature phenotypes through spatial-temporal incorporation of cell differentiation and chondrogenesis.

Keywords: Chondrogenic differentiation, chondrogenesis, mesenchymal stem cells, chondrocytes, chondrogenic differentiation factors.

Introduction

Articular cartilage is an avascular connective tissue with limited capacity for self-regeneration. Injury and degeneration of cartilage have far-reaching impacts on personal life, healthcare expenses and reduced productivity in the work force (Ge *et al.*, 2006). Current clinical treatment strategies, such as autologous chondrocyte implantation, bone marrow stimulation and mosaicplasty, have varying success rates, but their long-term results are far from satisfactory due to the lack of structural organisation of articular cartilage, as well as inferior mechanical properties of the newly formed tissue (Hunziker, 2009). Cartilage tissue engineering has the potential to create a more durable and functional replacement for the degenerated tissue; chondrocytes and stem cells are two promising cell source candidates for this replacement tissue. Chondrocytes are potentially the ideal seed cells with mature and stable phenotype as well as lineage stability. However, they are limited by scarcity during harvest and dedifferentiation during *in vitro* proliferation (Kock *et al.*, 2012). Establishment of embryonic stem cells (ESCs) requires embryos, which leads to ethical issues for clinical applications. Induced pluripotent stem cell (iPSC) technology can provide patient-specific cells, but uncontrolled and unexpected differentiation limits their application (Blin *et al.*, 2010; Yoshida and Yamanaka, 2010). Multipotent mesenchymal stromal cells or mesenchymal stem cells (MSCs) are a broadly adopted cell source for cartilage regeneration due to ample availability from multiple tissues, high proliferation capacity and the ability to differentiate into chondrocytes (Derfoul *et al.*, 2006; Kassis *et al.*, 2006). However, concomitant premature phenotypes of the differentiated chondrocytes, undesirable hypertrophic differentiation (Studer *et al.*, 2012), lack of subtype phenotypes and subsequent immature extracellular matrix (ECM) under current chondrogenic differentiation protocols have hindered the clinical application of stem cell-differentiated chondrocytes (Huey *et al.*, 2012; Mueller and Tuan, 2008).

Weighing the advantages and disadvantages of candidate cell sources, apart from using chondrocytes, MSCs may be the most adaptive cell source for current clinical use and for cartilage tissue engineering applications. Therefore, finding ways to improve *in vitro* chondrogenic differentiation of MSCs may be the next primary focus of tissue engineering research, which is somewhat different with the well-

* Address for correspondence:

Zigang Ge

Department of Biomedical Engineering, College of Engineering, Peking University, P.R China, 100871,

Telephone number: +86-10-62756736

Fax number: +86-10-62757545

Email: gez@pku.edu.cn

orchestrated embryonic chondrogenic differentiation process of MSCs. Embryonic chondrogenesis have been reviewed extensively with the focus on mesenchymal cell condensation, chondroprogenitor cells, chondrocytes as well as related mediators (Goldring *et al.*, 2006). Apart from embryonic chondrogenesis, joint specification and postnatal development have also been reviewed, though many questions remain unexplored (Onyekwelu *et al.*, 2009). MSCs have been well studied, regarding their definition, tissue origins, chemokines and receptors, as well as their *in vivo* microenvironments (Augello *et al.*, 2010); key factors like cell metabolic activity and transcript levels important for functional MSC-derived cartilage regeneration are summarised elsewhere (Chen *et al.*, 2006). Based on all these progresses, we summarise the history and challenges faced by current broadly used chondrogenic differentiation protocols with an aim to shed valuable light on the potential novel strategies to circumvent the challenges faced, which may benefit bioengineers and researchers in the field of tissue engineering and regenerative medicine.

Chondrogenic differentiation into articular chondrocytes

Chondrogenesis is initiated by cell condensation during embryonic development, before the mesenchymal stem cells differentiate into multiple lineages, such as articular, growth plate, intervertebral disc chondrocytes or endochondral ossification [some of the cells survive and transform into osteoblasts (Zhang *et al.*, 2014)]. For years, mesoderm was thought to be the origin of MSCs; however, new evidence indicates that the neuroepithelium, including the neural crest, may also be an early origin of MSCs (Morikawa *et al.*, 2009; Takashima *et al.*, 2007). MSCs from the somatopleure of the lateral plate mesoderm, neural crest cells in the neural ectoderm and sclerotome compartment of the paraxial mesoderm or somite differentiate into chondrogenic lineages and then yield the limbs, craniofacial bones and axial skeleton separately (Goldring *et al.*, 2006; Olsen *et al.*, 2000). Chondrogenesis may occur in one of two directions in the developing bone. One leads to the formation of bone *via* endochondral ossification, (at the primary and secondary centres of ossification, and in the growth plate) and the other leads to stable hyaline articular cartilage. The growth plate is composed of the resting zone, proliferative zone and hypertrophic zone (Abad *et al.*, 2002). Growth plate cartilage progresses to form the bone (Mackie *et al.*, 2008), whereas the articular cartilage does not (Pacifci *et al.*, 2005). Chondrocytes of the presumptive articular surface arise directly from a subpopulation of early chondroprogenitor cells, expressing doublecortin and growth/differentiation factor-5 (GDF-5) (Koyama *et al.*, 2008; Zhang *et al.*, 2007), but not matrilin-1 (Hyde *et al.*, 2007). Articular chondrocytes maintain the phenotype instead of resorting to hypertrophy by maintaining high expression of *ACAN* and *PRG4* (lubricin gene) and low expression of *COL2A1* with low proliferation (Lefebvre and Smits, 2005).

While articular cartilage at birth undergoes ossification and mineralisation, appositional growth from the surface forms an adult articular cartilage until puberty (Hunziker *et al.*, 2007; Onyekwelu *et al.*, 2009). These slowly dividing stem cells have bidirectional mitotic activity. In the horizontal direction, they furnish new stem cells that replenish the pool and contribute to the lateral expansion of the articular cartilage layer. In the vertical direction, the superficial zone supplies the rapidly dividing, transit-amplifying daughter-cell pool that feeds the transitional and upper radial zones and gradually forms the layered structure (Hunziker *et al.*, 2007). This growth activity of the articular surface ceases at puberty, whereas expansion of the metaphyseal growth plate continues until the time of skeletal maturity (Clark *et al.*, 1997; Hunziker *et al.*, 2007). Knowledge derived from endochondral ossification has been largely used to establish current chondrogenic differentiation protocols, while growth plate chondrocytes that are easily accessible sources are often compared to articular chondrocytes (Iwamoto *et al.*, 2013; Leijten *et al.*, 2012) (Fig. 1).

A history of the understanding of chondrogenic differentiation

Studies of the embryonic development of cartilage or cartilaginous tissues started as early as the 1960s. Ectopic implantation was a major research model during this period. The somites, which are embryonic cartilaginous progenitor cells, differentiated into chondrocytes *in vitro* when co-cultured with notochords (Lash *et al.*, 1960). Holtzer observed chondrogenesis of somites occurring in the presence of neural tubes (Holtzer, 1952; Holtzer and Detwiler, 1953). A fraction obtained from a cold perchloric acid extract of spinal cord and notochord was able to induce chondrogenesis (Lash *et al.*, 1962), which contained chondrogenic factors consisting of polypeptides, nucleotides and sugars (Holtzer, 1964; Hommes *et al.*, 1962). De-differentiation of chondrocytes was reported with morphological changes from spherical to stellate coupled with decreased synthesis of chondroitin sulphate and enhanced cell proliferation (Abbott and Holtzer, 1966). These findings led to the establishment of the pellet culture system, now broadly used in chondrogenic differentiation (Hattori and Ide, 1984).

In the 1970s and 1980s, laboratory studies evaluating the effect of cell shape, cell-cell and cell-matrix interactions on chondrogenesis were undertaken. Chondrocytes with spherical morphologies synthesised higher levels of chondroitin sulphate in comparison to those with spindle-like morphologies (Archer *et al.*, 1982). Cell-cell interactions, including both interactions amongst limb bud mesenchyme cells (Owens and Solursh, 1982) and epithelial-mesenchymal cells (Sanders, 1988), are essential for the differentiation of prechondrogenic limb mesenchyme into cartilage. Gene expression of both *COL1* and *fibronectin* increased at the onset of condensation, reached the highest point of chondrogenic differentiation and then subsided and were replaced with elevated *COL2A1* expression (Dessau *et al.*, 1980; Kulyk *et al.*,




Fig. 1. A schematic representation of different stages of chondrogenesis and endochondral ossification. The development of cartilage starts with MSC condensation. After committing to chondrogenesis, cells follow one of two different directions in the developing bone. One leads to the formation of bone *via* endochondral ossification (at the primary ossification centre (POC), secondary ossification centre (SOC), and in the growth plate) (the process marked in black) and the other that leads to stable hyaline articular cartilage. All committed hyaline chondrocytes start expressing *COL2A1*; matrilin-1 expression by cells arising from the interzones that are destined to become growth plate chondrocytes distinguishes them from articular chondrocytes. Doublecortin and GDF-5 only express in the articular cartilage. Most embryonic articular cartilage present at the ends of the bone at birth is replaced by bone and new cartilage formed by stem cells in the superficial zone. Articular cartilage becomes fully mature in adulthood. With the disappearance of growth plate cartilage, the SOC connects with the POC and gradually disappears in mature mammals. Stem-like cells exist in the cell condensation stage, keep stemness in the superficial zone of AC from birth until puberty, and exist in the form of resting stem-like cell after adulthood.

1989). Furthermore, in the 1980s, transforming growth factor- β (TGF- β) was identified as a key cartilage-inducing factor (Heine *et al.*, 1987; Seyedin *et al.*, 1986).

Many components play supplementary roles in the current chondrogenic differentiation medium. For instance, ascorbate stabilises the triple helical structures of collagen through hydroxylation of proline residues in procollagen and hydroxyproline (Peterkofsky, 1972). Serum inhibits the production of cartilage matrix during *in vitro* chondrogenesis of limb bud mesodermal cells of chick embryos (Hattori and Ide, 1984), whereas serum-free medium containing insulin/transferrin/selenium (ITS) successfully keeps cells alive and active (Bottenstein and Sato, 1979).

The current broadly-used chondrogenic differentiation medium was established in the 1990s, with the presence of TGF- β in pellet cultures (Johnstone *et al.*, 1998). This chondrogenic differentiation medium includes TGF- β , ITS, high-glucose, dexamethasone (Dex), ascorbic acid, sodium pyruvate and proline, but lacks serum (Johnstone *et al.*, 1998; Vater *et al.*, 2011) (Table 1). High-glucose was found to promote MSC survival and protect cells from apoptosis by altering the metabolism of the cells in pellet cultures and increasing the synthesis of proteoglycans through respiratory repression (Crabtree effect) (Mackay *et al.*, 1998).

The functions and mechanisms of differentiation-related growth factors, such as bone morphogenetic protein (BMP), fibroblast growth factor (FGF), parathyroid hormone related protein (PTHrP) and Indian hedgehog (Ihh), were studied in depth during this period (Murtaugh *et al.*, 1999; Shukunami *et al.*, 1996). Though these growth factors can benefit chondrogenesis *in vitro*, they are not routinely included in the current medium due to

several limitations. BMPs have been shown to promote the expression of *SOX9* and *COL II*, however, they also inevitably result in a hypertrophic phenotype of cells (Enomoto-Iwamoto *et al.*, 1998; Nonaka *et al.*, 1999). FGFs can maintain the proliferation of cells but result in chondrodysplasia when overexpressed (Minowada *et al.*, 1999). Additionally, PTHrP can reverse chondrocytes from a hypertrophic phenotype to a prehypertrophic proliferating phenotype, and prevent terminal differentiation of chondrocytes *in vitro* (Zerega *et al.*, 1999). Ihh expressed in prehypertrophic chondrocytes enhances hypertrophic differentiation. Misexpression of Ihh can prevent hypertrophic differentiation of chondrocytes (Vortkamp A, 1996).

In the late 1990s, the role of integrins in mediating cell attachment to extracellular matrix proteins was highlighted. These interactions regulated morphogenesis and cell differentiation. Treatment with integrin $\beta 1$ antibodies inhibited early chondrogenesis of limb bud cells from mouse embryos *in vitro* (Shakibaei, 1998). Additionally, *SOX9* was found to be a master gene which regulated downstream *COL2A1* expression directly (Healy *et al.*, 1999).

Understanding of chondrogenic differentiation has accelerated since the year 2000, attributed to the application of a variety of genetic modification techniques. Epigenetics then emerged as an important field. MicroRNAs (miRNAs) proved effective mediators of key pathways during MSC differentiation through regulating transcription factors (Hong and Reddi, 2012). DNA methylation and histone modification (including histone acetylation and deacetylation) are involved in chondrogenesis through modulating chromatin structures, which directly influence the activity of DNA in transcription, replication and

Table 1. Chondrogenic differentiation medium and effects of individual components on chondrogenesis

	Factors	Function	Signalling	Dose	Synergy	Subtypes
The well-established chondrogenic differentiation serum-free media (Johnstone <i>et al.</i> , 1998; Vater <i>et al.</i> , 2011)	TGF-β (J.Zuscik <i>et al.</i> , 2004; Lutz and Knaus, 2002; Park <i>et al.</i> , 2011; Schmierer and Hill, 2007)	Promotes chondrogenesis, enhances extracellular matrix production and downregulates collagen I expression	MAPK, Wnt, β -catenin, Smad	10 ng/mL	IGF-I, BMPs, PHTrP	TGF- β 1, β 2 and β 3, have similar functions
	Insulin (Mueller <i>et al.</i> , 2013; Quarto <i>et al.</i> , 1992)	Stimulates collagen synthesis, inhibits hypertrophy	Enhances tyrosine kinase activity of the insulin receptor	1-10 μ g/mL	Transferrin, selenium	
	Transferrin (Cigan, 2013; Kisiday <i>et al.</i> , 2005)	Iron-binding		1-30 μ g/mL	Insulin, selenium	
	Selenium (Yan <i>et al.</i> , 2012)	Anti-oxidant and enhances proliferation		5 μ g/mL	Insulin, transferrin	
	Dex (M.Florine <i>et al.</i> , 2013; Thomas M.Randua, 2013)	Supports cell viability and delays the appearance of type X collagen	GRs, Scrapie Responsive Gene 1	10 ⁻⁷ M	TGF- β	
	Glucose (Cigan, 2013; Han <i>et al.</i> , 2004; Tsai, 2013)	Major energy source, precursor for the synthesis of glycosaminoglycans	PKC α , p38MAPK, ERK	4.5 g/L		
	Ascorbic acid (Choi <i>et al.</i> , 2008; Franceschi, 1992; Peterkofsky, 1972; Stone and Meister, 1962)	Proliferation, DNA synthesis, collagen biosynthesis and reductant	Reduction of iron to its ferrous state	50 mg/mL	Proline	L-ascorbate, sodium L-ascorbate, and L-ascorbate-2-phosphate
	Pyruvate (Andrae <i>et al.</i> , 1985; Geshi <i>et al.</i> , 2000)	Energy source, against hydrogen peroxide-induced cytotoxicity		1 mM		
	Proline (Peterkofsky, 1972; Washington <i>et al.</i> , 2010)	Stabilisation of the collagen triple helix, promotes ES cell differentiation	mTOR signalling pathway	40 μ g/mL	Ascorbic acid	
Selected growth factors regulators	BMPs (Cao and Chen, 2005; Hassel <i>et al.</i> , 2003; Hatakeyama <i>et al.</i> , 2004; Jin <i>et al.</i> , 2006; Sekiya <i>et al.</i> , 2005; Shen <i>et al.</i> , 2010)	Enhances cartilage formation	MAPK, Wnt, Smad	10-500 ng/mL	TGF- β	BMP-2, enhances proliferation and ECM, and downregulates Col I expression. BMP-4, accelerates the progression of cartilage differentiation to maturation BMP-6, enhances chondrogenesis in special subpopulation of MSCs, increases proteoglycans BMP-7, inhibits cell proliferation, induces chondrogenic differentiation BMP-9, enhances Sox9, Col2A1 and aggrecan, overcome the inhibitory effect of IL-1. GDF-6(BMP-13), inhibits hypertrophic chondrocyte specific marker, upregulate of proteoglycan
	GDF-5(BMP-14) (Coleman and Tuan, 2003; Hatakeyama <i>et al.</i> , 2004)	Enhances cell condensation, chondrogenesis, and ECM production	Gap junction intercellular communication, p38MAPK	150 ng/mL	TGF- β , BMP	
	PHTrP (Barbara Zerega, 1999; Kim <i>et al.</i> , 2008; Mau <i>et al.</i> , 2007)	Inhibits the TGF- β -induced hypertrophic differentiation	PTHrP and IHH feedback loop	10 μ M	TGF- β	PHTrP1-34
	FGF (Ng <i>et al.</i> , 2008; Solchaga <i>et al.</i> , 2005; Tsutsumi <i>et al.</i> , 2001)	Regulates proliferation, and increases cartilaginous ECM production	ERK	10 ng/mL FGF-2 10 ⁻⁹ M FGF-18	TGF- β	FGF-2, increases proliferation, and proteoglycan production FGF-18, inhibits cell proliferation and induces chondrogenic differentiation

recombination (Furumatsu and Ozaki, 2010; Roth *et al.*, 2001). Recently, several groups found an important relationship between epigenetic marking and gene expression in chondrogenesis. Herlofsen *et al.* found that modifications can provide primary epigenetic control of the early differentiation of MSCs toward the chondrogenic lineage (Herlofsen *et al.*, 2013).

Biophysical, chemical and mechanical factors were found to be heavily involved in chondrogenesis (Leijten *et al.*, 2014). Small chemical molecules (Benoit *et al.*, 2008), matrix elasticity (Engler *et al.*, 2006; Park *et al.*, 2011), cell material property (cell softness) (Chowdhury *et al.*, 2010) and many more factors, are now known to guide MSCs to differentiate into specific cell lineages. A breakthrough

in the regeneration of articular cartilage illustrated that an entire articular surface can be regenerated using adult stem/progenitor cells recruited from the host with the help of biomaterial-based scaffold and growth factors (Lee *et al.*, 2010).

Key elements in current chondrogenic differentiation medium

At present, MSCs cultured in pellet and micromass systems in the presence of chondrogenic differentiation medium are the two major methods for initiating chondrogenic differentiation *in vitro*. The following sections will describe




Fig. 2. TGF- β and MAPK signalling in chondrogenic differentiation of MSCs.

the in-depth role of the ingredients potentially used in chondrogenic differentiation medium.

TGF- β is the core component in most chondrogenic differentiation protocols used currently. TGF- β s exist in three homologous homodimeric forms (β 1, β 2 and β 3) and all of them signal through the same receptors [TGF- β receptor I (TGF- β RI) and II (TGF- β RII)]. After activation of the TGF- β receptors, TGF- β induces chondrogenesis mainly through TGF- β /drosophila mothers against decapentaplegic protein (Smad) and mitogen-activated protein kinase (MAPK) signalling pathways, that cross-talk with each other (Li *et al.*, 2010; Lutz and Knaus, 2002; Schmierer and Hill, 2007) (Fig. 2). Before phosphorylation by TGF- β RII, TGF- β RI is catalytically inactive due to a wedge-shaped GS region (a highly conserved 30-amino acid region with a characteristic SGSGSG sequence) inserted into the kinase domain (Massagué, 1998). In Smad signalling, TGF- β RII phosphorylates the GS region after forming a ligand-receptor complex with TGF- β , which results in the activation of TGF- β RI (Huse *et al.*, 1999). The activated TGF- β RI phosphorylates Smad2/3, which recruits Smad4, and the complex translocates into the nucleus (Massagué and Wotton, 2000). The translocated Smad2/3 associates with Sox9 and CBP/p300 (cAMP-response element-binding protein, which has an intrinsic histone acetyltransferase activity and acts as an important co-activator for the expression of the *COL2A1*) on the *COL II* enhancer region containing the Sox9-binding site that is important for the initiation of chondrogenic differentiation (Furumatsu *et al.*, 2005). As for MAPK

signalling, the major subtypes of MAPK include p38, extracellular signal-regulated kinase-1 (ERK-1) and c-jun N-terminal kinase (JNK). In addition, p38 is an enhancer of chondrogenesis, whereas ERK-1 is a repressor of chondrogenesis; JNK plays a minor role in chondrogenesis. Activation of MAPK involves indirect modulation of cell adhesion molecules, including N-cadherin and integrin $\alpha_5\beta_1$, during precartilaginous condensation and progression to chondrogenic differentiation (Tuli *et al.*, 2003). TGF- β induces the expression of the chondrogenic factor *SOX9* and the function of the main chondrogenic trio (*SOX5*, *SOX6* and *SOX9*) (Lefebvre *et al.*, 1998; Lefebvre and Smits, 2005) that further regulates chondrogenic differentiation and ECM formation. It is known that the genes for many ECM components, such as *ACAN*, *COL2A1* and *Matrilin-1*, are only expressed when the three Sox proteins bind to the high-mobility group (HMG)-domain sites in their enhancers (Akiyama and Lefebvre, 2011). On the contrary, TGF- β also plays an inhibitory role in chondrocyte maturation (Ballock *et al.*, 1993; Kato *et al.*, 1988; Serra *et al.*, 1997).

ITS premix is a substitute for foetal bovine serum that maintains cellular activity in biosynthesis and cell division. Serum has some disadvantages including poor characterisation and inconsistency in batch-to-batch composition and quality (Barnes and Sato, 1980; Cigan, 2013). Insulin is a primary factor involved in the onset and progression of chondrogenesis (Quarto *et al.*, 1992) and can facilitate chondrogenesis in a dose-dependent manner (1-10 μ g/mL) with added TGF- β (Mueller *et al.*,

2013a). Furthermore, insulin can facilitate glucose uptake (Kono *et al.*, 1982) and enhance DNA synthesis and proteoglycan production in cartilage cultures (Maor *et al.*, 1993; Rosen, 1987). Transferrin promotes cell proliferation and differentiation through detoxifying oxygen radicals and peroxides (Ned *et al.*, 2003; Schäfer *et al.*, 2007). Selenite, a co-factor for glutathione peroxidase and other proteins, enhances proliferation of chondrocytes through acceleration of G1-phase cell cycle progression and the induction of cyclin D1 (Wei *et al.*, 1986). This effect is mediated by the enhancement of intracellular adenosine triphosphate (ATP) content (Yan *et al.*, 2012).

Dex, a synthetic glucocorticoid, promotes chondrogenic differentiation through enhancing related gene expression (Florine *et al.*, 2013; Jakobsen *et al.*, 2014; Randua *et al.*, 2013). The chondrogenic effects of glucocorticoids are predominantly mediated *via* the glucocorticoid receptors (GRs). GR α constitutes the main active GR form while GR β constitutes a negative isoform of GR α (Oakley *et al.*, 1999). Dex promotes chondrogenic differentiation by down-regulating GR β levels and maintaining GR α levels (Derfoul *et al.*, 2006). Dex enhances the gene expression of *COL XI* under chondrogenic conditions (collagen XI acts as both the template for collagen II fibrillogenesis and a regulator for maintaining fibril diameters in cartilage) (Chen *et al.*, 2005; Derfoul *et al.*, 2006). Previous studies have shown that Dex enhanced the TGF- β induced expression of *ACAN*, *COL2A1* and cartilage oligomeric matrix protein (*COMP*) and supported cell viability, as well as delayed the expression of *COL X* when used in conjunction with TGF- β (Derfoul *et al.*, 2006; Quarto *et al.*, 1992). The tissue origin of MSCs determines the effect of Dex on chondrogenic differentiation. Dex enhanced TGF- β 1-induced chondrogenesis in bone marrow-derived MSCs (BMSCs) but had no significant impact on the TGF- β 1- or BMP2-induced response in synovial derived MSCs (Kurth *et al.*, 2007; Park *et al.*, 2005; Shintani and Hunziker, 2011; Shintani *et al.*, 2007). This may be due to the different cell survival rates and microenvironment which could be tissue-source dependent (Buxton *et al.*, 2011).

Glucose, a major energy source and precursor of ECM for most mammalian cells, promotes chondrogenic differentiation and inhibits apoptosis at high levels (4.5 g/L or 25 mM) (Mackay *et al.*, 1998; Shikhman *et al.*, 2001; Tsai *et al.*, 2013). Glucose participates in multiple metabolic cycles of MSCs during chondrogenic differentiation, including oxygen consumption, glucose consumption and lactate production. High levels of glucose in the medium enhance cell differentiation by changing the metabolic patterns and modulating subsequent cell signalling pathways. The synthesis of proteoglycans increases through progressively inhibiting the respiration of chondrocytes cultured *in vitro*, as the glucose concentrations (1-10 mM) increase (Derfoul *et al.*, 2006; Otte, 1991). High doses of glucose enhance the differentiation of stem cells into chondrogenic lineages through the down-regulation of ERK and protein kinase C α (PKC α) activities, up-regulation of p38, as well as through modulating the expression of adhesion molecules such as fibronectin, integrin β 1 and N-cadherin (Han *et al.*, 2004).

Both ascorbic acid and proline are essential for the

production of collagen in cartilage. Ascorbic acid enhances cell proliferation, synthesis and extracellular deposition of collagenous matrix, in particular collagen I and II (Choi *et al.*, 2008; Franceschi, 1992; Potdar and D'Souza, 2010; Stone and Meister, 1962). Ascorbic acid stabilises the collagen triple helical structure through reduction of iron to its ferrous state, which is important for the hydroxylation of proline residues in procollagen and hydroxyproline and the maturation of procollagen to collagen (Padh, 1991; Peterkofsky, 1991). Proline has been shown to regulate ESC differentiation into early primitive ectoderm-like (EPL) cells, through mammalian targeting of the rapamycin (mTOR) signalling pathway (Washington *et al.*, 2010).

Sodium pyruvate is essential for enhancing the energy metabolism in the Krebs cycle (Geshi *et al.*, 2000). Mammalian cells secrete pyruvate and α -ketoacids in culture to protect them against hydrogen peroxide-induced cytotoxicity (Andrae *et al.*, 1985; O'Donnell-Tormey *et al.*, 1987). Pyruvate up-regulates the expression of genes involved in free radical scavenging which, in turn, enhances the antioxidative machinery and is essential for maintaining mitochondrial activity. Super-physiological concentrations of pyruvate (50 mM, 50 folds of physiological concentrations) enhance mitochondrial mass and functionality (Wilson *et al.*, 2007).

Phenol red, a commonly used pH indicator in tissue culture medium can inhibit the chondrogenic differentiation of MSCs (Lysdahl *et al.*, 2013). During chondrogenic differentiation, phenol red in Dulbecco's Modified Eagle Medium (DMEM) has been shown to decrease *SOX9*, *COL2A1* and *ACAN* on days 14 and 21 and proteoglycan synthesis on days 21 and 28 in culture (Lysdahl *et al.*, 2013). This phenomenon may have occurred due to phenol red's structural resemblance to certain nonsteroidal oestrogens and weak oestrogen agonist activity (Berthois *et al.*, 1986).

BMPs are morphogens in the TGF- β superfamily and regulate chondrogenesis and skeletogenesis during normal embryonic development (Hogan, 1996). They are involved in various stages of chondrogenic differentiation, from initiation of chondrogenic differentiation, to regulation of chondrocyte maturation and terminal differentiation (Pizette and Niswander, 2000). BMP2, BMP4, BMP6 and BMP7 are the most commonly used BMPs for chondrogenic differentiation (Kramer *et al.*, 2000; Sekiya *et al.*, 2005; Shen *et al.*, 2010). BMPs signal through Smad or non-Smad signalling pathways. The Smad1/5/8 route is an important Smad-dependent pathway that induces hypertrophy and bone formation (Cao and Chen, 2005). The alternative Smad-independent pathway in BMP regulates chondrogenesis, mediated by activating several MAPKs including ERKs and p38 kinases (Hassel *et al.*, 2003). p38 enhances chondrogenesis by up-regulating *SOX9* expression directly or by inhibiting the Wnt7a/beta catenin pathway indirectly (Jin *et al.*, 2006). MSCs from different species and tissue sources vary in their responsiveness to BMPs, partly due to their distinct cell receptor repertoires (Hennig *et al.*, 2007; Osyczka *et al.*, 2004).

GDF5, also known as BMP14 in humans, is a member of the TGF- β superfamily and plays an important

role in limb bud mesenchymal cell condensation and chondrogenesis during embryogenesis (Coleman and Tuan, 2003; Storm *et al.*, 1994). GDF5 stimulates the recruitment and differentiation of chondrogenic cells at early stages and increases chondroprogenitor cell condensation and cartilaginous nodules without altering the overall pattern of differentiation. It causes a more sustained elevated expression level of *SOX9* than *BMP4* (Hatakeyama *et al.*, 2004). GDF5 increases mesenchymal cell condensation independent of cell density or N-cadherin-mediated adhesion and signalling; however, it is dependent on gap junction-mediated cellular communication (Coleman and Tuan, 2003; Sun *et al.*, 2012). GDF5 can promote chondrogenesis *via* signalling cross-talk, when used together with TGF- β 3 and BMP2 *in vitro* (Murphy *et al.*, 2015). GDF5 has also been shown to increase the expression of *COL2A1* and hypertrophic marker, alkaline phosphatase (*ALP*) by enhancing the phosphorylation of Smad1/5/8 (Coleman *et al.*, 2013), which is activated by BMPs. GDF5's effect on the enhancement of *ACAN* and the inhibition of *MMP13* expression occurs by up-regulating the canonical Wnt inhibitors DKK1 (Enochson *et al.*, 2014).

PTHrP inhibits TGF- β -induced hypertrophic differentiation by up-regulating the chondrogenic markers *COL2A1* and *SOX9* and down-regulating the hypertrophic markers *COL10A1* and *RUNX2* during *in vitro* chondrogenesis of MSCs (Kim *et al.*, 2008; Weiss *et al.*, 2010). PTHrP exerts its effects *in vivo* *via* the PTHrP and IHH feedback loop by binding to its receptor (PTHR1) and activating a range of signalling molecules, *e.g.* protein kinase A (PKA), PKC and inositol 1, 4, 5-tris-phosphate (IP3). Some signalling molecules act as negative regulators for IHH. Hence, PTHrP inhibits the terminal differentiation of chondrocytes driven by IHH (Kim *et al.*, 2008; Mau *et al.*, 2007; Rabie *et al.*, 2003). The chondrogenic effects of PTHrP vary amongst different PTHrP isoforms. PTHrP 1-34, added in chondrogenic medium containing TGF- β 3, significantly increases chondrogenic differentiation with less hypertrophic differentiation, while other isoforms (PTHrP 1-86, 7-34 and 107-139) show inconsistent effects (Lee and Im, 2012). When culturing MSC pellets under hypertrophy-enhancing conditions [hypertrophy-enhancing medium consisting of DMEM, 1 % ITS, 50 μ g/mL ascorbate-2-phosphate, 40 μ g/mL L-proline and 1 nM triiodothyronine (T3)], adding PTHrP (1-40) does not diminish the induced enhancement of hypertrophy (Mueller *et al.*, 2013b).

The FGF family, especially FGF [basic FGF (bFGF); also known as FGF2], can maintain MSCs in an immature state during *in vitro* expansion and enhance MSC proliferation and differentiation potential (Mastrogiacomo *et al.*, 2001; Solchaga *et al.*, 2005; Tsutsumi *et al.*, 2001). Literature suggests that aggregates of human MSCs that were expanded in culture in the presence of bFGF lacked the collagen I-positive and collagen II-negative outer layer characteristics of aggregates (Ng *et al.*, 2008; Solchaga *et al.*, 2005). However, in pellet cultures, the prolonged exposure of MSCs to bFGF caused the reduction of glycosaminoglycan (GAG) production (Hellingman *et al.*, 2010). Therefore, bFGFs can be added into the medium to

culture MSCs, but in differentiation medium, it may not be necessary.

Physical, chemical and biological factors

In addition to biochemical molecules used in current chondrogenic differentiation medium, many other factors are involved in chondrogenic differentiation. These factors include physical (oxygen tension, mechanical, topography, *etc.*), chemical (small molecules, chemical properties, *etc.*) and biological factors (gene modification, bioactivity, epigenetics, *etc.*). Apart from biochemical factors (growth factors), biophysical factors (co-culture) and biophysicochemical factors (biomaterials) are equally important (Fig. 3). The influence of biophysical factors on chondrogenesis *in vivo* and *in vitro* have been reviewed (Huang *et al.*, 2010a; Studer *et al.*, 2012; Tate *et al.*, 2008). In this section, we focus on the use of biophysical and biochemical cues for chondrogenesis and recent advances in the field of tissue engineering.

Hypoxia can facilitate chondrogenic differentiation of MSCs, in addition to maintaining an undifferentiated state and inducing apoptosis. The microenvironment or the niche of MSCs in the bone marrow has low oxygen tension (< 32 mm Hg, 4.2 % oxygen) (Oze *et al.*, 2012; Spencer *et al.*, 2014). Therefore, chondrogenic differentiation is enhanced when MSCs are cultured in hypoxia (5 % oxygen) in comparison to normoxia (20 % oxygen) (Lee *et al.*, 2013). Hypoxia has a positive influence on chondrogenic differentiation of MSCs, mainly through phosphorylation of AKT (known as protein kinase B) and p38 MAPK, thereby down-regulating hypoxia-inducible factor (HIF-1) (Kanichai *et al.*, 2008).

Mechanical stimulation can directly influence the fate of undifferentiated stem cells (Estes *et al.*, 2004) *via* cellular and nuclear deformation and indirect biophysical factors, including osmotic and hydrostatic pressure and fluid flow (O'Connor *et al.*, 2013). Matrix elasticity directs the differentiation of stem cells; soft matrix (~ 1 kPa) has been shown to promote MSC differentiation into a chondrogenic lineage (Engler *et al.*, 2006; Park *et al.*, 2011), while activation and internalisation of integrin are actively involved in the regulation of stem cell differentiation (Du *et al.*, 2011). Cell-material properties, such as cell stiffness, dictate cell spreading and alter differentiation in response to physical stimuli (Chowdhury *et al.*, 2010). Surface topography can trigger changes in MSC morphology, gene expression and cytoskeletal structure (Huang *et al.*, 2010b). Compared to non-patterned surfaces, nano-pillar and nano-hole topographies enhanced MSC chondrogenesis and facilitated hyaline cartilage formation, whereas nano-grill topography delayed chondrogenesis and induced the formation of fibro/superficial zone cartilage (Wu *et al.*, 2014).

Small chemical molecules are also known to guide MSC differentiation (Benoit *et al.*, 2008) by binding to specific receptors that promote chondrogenic differentiation (Cho *et al.*, 2012; Johnson *et al.*, 2012; Kafienah *et al.*, 2007; Schugar *et al.*, 2008). As an illustrative example, kartogenin (KGN) has been shown to encourage




Fig. 3. Factors in the process of *in vitro* chondrogenic differentiation in MSCs.

chondrogenic differentiation of human MSCs, by significantly enhancing the expression of *COL2A1*, *SOX9* and *ACAN*. However, KGN had no significant effect on the gene products associated with chondrocyte calcification, such as osteocalcin (*OCN*), *ALP* and *COL10A1*, in human MSCs or chondrocytes (Johnson *et al.*, 2012). KGN functions by binding to filamin A, an actin-binding protein that cross-links actin filaments. Thereby, by regulating the cytoskeletal network organisation and dynamics (Gorlin *et al.*, 1990) and disrupting its interaction with core binding factor β (CBF β), it modulates the activity of the *RUNX* family of transcription factors (Johnson *et al.*, 2012).

Small molecule inhibitors can regulate chondrogenesis by blocking key receptors or signalling pathways. As a nuclear receptor, retinoic acid receptor β (RAR β) is down-regulated significantly during chondrogenesis. The synthetic inhibitor (LE135) enhances chondrogenesis while inhibiting hypertrophic differentiation (Kafienah *et al.*, 2007). Another small molecule inhibitor, dorsomorphin, can block Smad1/5/8 signalling, enhance chondrogenesis, elevate the production of collagen II and prevent mineralisation (Hellingman *et al.*, 2011; Retting *et al.*, 2009). Inhibition of inflammatory factors can boost the chondrogenic differentiation capacity of MSCs obtained from osteoarthritis patients through inhibiting inflammatory stimuli-induced damage, hypertrophy and apoptosis (Murphy *et al.*, 2002; Rainbow *et al.*, 2013).

Using a gene transfer/silencing strategy on MSCs can facilitate change in the complex regulatory networks of stem cell biology and regulate chondrogenic differentiation processes (Feng *et al.*, 2008; Paik *et al.*, 2012). For example,

knockdown of HIF-1 α by HIF-1 α small interfering RNA (siRNA) leads to an increase in chondrogenesis (Kanichai *et al.*, 2008).

Biomaterials can facilitate chondrogenesis in multiple ways, including mechanical support, surface topography and stiffness, chemical modification and bioactivity (Bian *et al.*, 2013; Lutolf *et al.*, 2009; Wise *et al.*, 2009; Zhang *et al.*, 2014). Co-cultures of chondrocytes and MSCs enhance chondrogenesis due to the trophic role of MSCs (MSCs secrete a variety of cytokines and growth factors that have both paracrine and autocrine activities), which stimulate chondrocyte proliferation and matrix production. However, MSCs in co-culture conditions do not actively undergo chondrogenic differentiation themselves (Caplan and Dennis, 2006; Meretoja *et al.*, 2012; Wu *et al.*, 2012).

Future Perspectives

Heterogeneous differentiation of differentiated MSCs is one of the main challenges faced in *in vitro* chondrogenic induction and is seldom discussed. Multiple factors are involved in these processes, such as variance in cell-cell and cell-matrix adhesions, autocrine or paracrine activities, different subpopulations of MSCs used, individual cells encapsulated in biomaterials that induces different cellular microenvironments (such as exchange of nutrient and waste, oxygen gradients), *etc.*. High-throughput technologies, such as microfluidics or microfabrication, may be able to set up large scale production of homogeneous microtissues or microenvironments to circumvent this issue

(Jakobsen *et al.*, 2014). Assembling pellets containing varying sub-phenotypes of differentiated chondrocytes layer-by-layer has previously been shown to mimic the structural properties of articular cartilage (Chen *et al.*, 2006). Defining subpopulations of MSCs may provide a good foundation; some progress in this field has been made (Houlihan *et al.*, 2012). As cells only differentiate when they exit the cell cycle, fine-tuned cell cycles at the initial stages of differentiation can influence cells to differentiate into homogeneous phenotypes (Li *et al.*, 2014).

Senescence and apoptosis inherently occur during both embryonic development and *in vitro* chondrogenic differentiation, so regulation of these two processes could potentially regulate chondrogenic differentiation. Senescence and apoptosis synergistically prevent the over-growth of chondrocytes and shapes the articular cartilage during embryonic development (Ito and Kida, 2000; Loeser, 2009). Senescence of bone marrow MSCs is associated with low proliferation activity and decline of stemness (Squillaro *et al.*, 2015), which may lead to inferior chondrogenic differentiation efficiency (Vacanti *et al.*, 2005). Efforts in hampering cell senescence during chondrogenic differentiation with growth factors, anti-inflammatory drugs, antioxidants, nutrients as well as appropriate microenvironments such as hypoxia and stem cell matrices effectively improve the efficiency and quality of tissue-engineered cartilage *in vitro* and *in vivo* (Li and Pei, 2012; Rainbow *et al.*, 2013). It is known that the majority of cells in pellet cultures undergo apoptosis (Wang *et al.*, 2010). Apoptosis may occur due to low nutrient supply to cells in the centre of the pellet, hypoxia and subsequent cell fate. Some preconditioning steps, such as use of medium with high glucose, and hypoxia preconditioning, may be able to protect MSCs from undergoing the process of apoptosis (Mackay *et al.*, 1998; Wang *et al.*, 2008).

In addition, biophysical signals are indispensable for successful chondrogenic differentiation. Mechanical properties influence phenotypes of differentiated chondrocytes and affect respective progenies (Yang *et al.*, 2014). Oxygen tension or hypoxia can facilitate differentiation. MSCs can differentiate into hyaline cartilage chondrocytes instead of hypertrophic chondrocytes which is more likely to happen under current chondrogenic differentiation protocols (Leijten *et al.*, 2014). Premature and unstable phenotypes of differentiated chondrocytes pose challenges for potential clinical use and lead to inferior quality of regenerated cartilage (Peltari *et al.*, 2006). Mimicking embryonic articular cartilage formation instead of embryonic chondrogenesis can assist bioengineers in overcoming this challenge, which means that temporal regulation of multiple factors may be required. For example, decreased Wnt, hedgehog and BMP signalling and specification of GDF-5 expression are of great importance for the homeostasis of articular chondrocytes (Bobacz *et al.*, 2008; Leijten *et al.*, 2012). Previous studies have shown that over expression of GDF-5 led to the unsuccessful differentiation of articular chondrocytes (Feng *et al.*, 2008). However, the role of GDF-5 in embryonic arthrogenesis and postnatal development is not entirely clear. From our viewpoint, hypertrophy

will not pose challenges to chondrogenic differentiation if optimised protocols can further differentiate stem cells into articular chondrocytes and maintain their classic phenotypes.

Temporally fine-tuning chondrogenic differentiation medium can profoundly enhance differentiation efficiency. Chondrogenic differentiation medium is involved in administering TGF- β and other factors throughout the entire culture period (Johnstone *et al.*, 1998). Previously, multiple factors induced step-wise mesoderm differentiation and subsequent chondrogenesis through mimicking embryonic chondrogenesis (Oldershaw *et al.*, 2010). The sequential addition of growth factors may be able to improve chondrogenesis based on gene expression of cell surface receptors and effectiveness of growth factors; withdrawal of the growth factors after nine days improves chondrogenesis (Handorf and Li, 2014). Apart from the addition of growth factors, inhibition of specific cell signalling pathways may also enhance chondrogenesis, such as the Smad1/5/8 signalling inhibitor dorsomorphin (Hellingman *et al.*, 2011; Retting *et al.*, 2009) or Wnt inhibitors [Gremlin 1, frizzled-related protein (FRP) or dickkopf (Dkk-1)] (Leijten *et al.*, 2012).

Use of mathematical models can assist researchers in regulating chondrogenic differentiation. A “seesaw model” describes a balance between mesendodermal and ectodermal specification of pluripotency (Shu and Deng, 2013) and the cross-talk between genetics and epigenetics (Leijten *et al.*, 2014). The lineage specifiers of mesendodermal and ectodermal are considered as pluripotency rivals. However, these lineage specifiers have shown to facilitate reprogramming of differentiated cells to the pluripotent state when a balance is achieved.

Summary

MSCs are a promising source of precursor cells, which are widely used in cartilage tissue engineering. By reviewing the process of cartilage development and the constituents of existing medium formulations for promoting chondrogenic differentiation, we have identified potential factors to generate a more stable and reliable hyaline cartilage *in vitro*. Cellular metabolic status, especially glucose metabolism and inhibition of inflammatory factors, need to be considered when trying to inhibit cell senescence and apoptosis. The combination and concentration of growth factors, the timing of the application of biochemical components and proper biophysical stimuli during cell culture needs further optimisation to achieve desirable differentiation outcomes.

Acknowledgements

The authors would like to thank Professor MB Goldring at Weill Cornell Medical College for her valuable advice on chondrogenesis and assistance with in editing this manuscript. We would also like to thank Ms. Suzanne Danley at West Virginia University and Dr. Ayeesha Mujeeb for proofreading this manuscript. The authors would like

to acknowledge support from the National Basic Research Program of China (973 Program) (2012CB619100) and National Natural Science Foundation of China grant (81271722, 81471800).

Disclosure Statement

No competing financial interests exist.

References

- Abad V, Meyers JL, Weise M, Gafni RI, Barnes KM, Nilsson O, Bacher JD, Baron J (2002) The role of the resting zone in growth plate chondrogenesis. *Endocrinology* **143**: 1851-1857.
- Abbott J, Holtzer H (1966) The loss of phenotypic traits by differentiated cells. *J Cell Biol* **28**: 473-487.
- Akiyama H, Lefebvre V (2011) Unraveling the transcriptional regulatory machinery in chondrogenesis. *J Bone Miner Metab* **29**: 390-395.
- Andrae U, Singh J, Ziegler-Skylakakis K (1985) Pyruvate and related alpha-ketoacids protect mammalian cells in culture against hydrogen peroxide-induced cytotoxicity. *Toxicol Lett* **28**: 93-98.
- Archer CW, Rooney P, Wolpert L (1982) Cell shape and cartilage differentiation of early chick limb bud cells in culture. *Cell Differ* **11**: 245-251.
- Augello A, Kurth TB, De Bari C (2010) Mesenchymal stem cells: a perspective from *in vitro* cultures to *in vivo* migration and niches. *Eur Cell Mater* **20**: 121-133.
- Ballock RT, Heydemann A, Wakefield LM, Flanders KC, Roberts AB, Sporn MB (1993) TGF-beta-1 prevents hypertrophy of epiphyseal chondrocytes - regulation of gene-expression for cartilage matrix proteins and metalloproteases. *Dev Biol* **158**: 414-429.
- Barnes D, Sato G (1980) Methods for growth of cultured cells in serum-free medium. *Anal Biochem* **102**: 255-270.
- Benoit DS, Schwartz MP, Durney AR, Anseth KS (2008) Small functional groups for controlled differentiation of hydrogel-encapsulated human mesenchymal stem cells. *Nat Mater* **7**: 816-823.
- Berthois Y, Katzenellenbogen JA, Katzenellenbogen BS (1986) Phenol red in tissue culture media is a weak estrogen: implications concerning the study of estrogen-responsive cells in culture. *Proc Natl Acad Sci USA* **83**: 2496-2500.
- Bian L, Guvendiren M, Mauck RL, Burdick JA (2013) Hydrogels that mimic developmentally relevant matrix and N-cadherin interactions enhance MSC chondrogenesis. *Proc Natl Acad Sci U S A* **110**: 10117-10122.
- Blin G, Nury D, Stefanovic S, Neri T, Guillevic O, Brinon B, Bellamy V, Rucker-Martin C, Barbry P, Bel A, Bruneval P, Cowan C, Pouly J, Mitalipov S, Gouadon E, Binder P, Hagege A, Desnos M, Renaud JF, Menasche P, Puceat M (2010) A purified population of multipotent cardiovascular progenitors derived from primate pluripotent stem cells engrafts in postmyocardial infarcted nonhuman primates. *J Clin Invest* **120**: 1125-1139.
- Bobacz K, Sunk I-G, Hayes S, Amoyo L, Tohidast-Akrad M, Kollias G, Smolen JS, Schett G (2008) Differentially regulated expression of growth differentiation factor 5 and bone morphogenetic protein 7 in articular cartilage and synovium in murine chronic arthritis: potential importance for cartilage breakdown and synovial hypertrophy. *Arthritis Rheum* **58**: 109-118.
- Bottenstein JE, Sato GH (1979) Growth of a rat neuroblastoma cell line in serum-free supplemented medium. *Proc Natl Acad Sci USA* **76**: 514-517.
- Buxton AN, Bahney CS, Yoo JU, Johnstone B (2011) Temporal exposure to chondrogenic factors modulates human mesenchymal stem cell chondrogenesis in hydrogels. *Tissue Eng Part A* **17**: 371-380.
- Cao X, Chen D (2005) The BMP signaling and *in vivo* bone formation. *Gene* **357**: 1-8.
- Caplan AI, Dennis JE (2006) Mesenchymal stem cells as trophic mediators. *J Cell Biochem* **98**: 1076-1084.
- Chen CW, Tsai YH, Deng WP, Shih SN, Fang CL, Burch JG, Chen WH, Lai WF (2005) Type I and II collagen regulation of chondrogenic differentiation by mesenchymal progenitor cells. *J Orthop Res* **23**: 446-453.
- Chen J, Horan RL, Bramono D, Moreau JE, Wang Y, Geuss LR, Collette AL, Volloch V, Altman GH (2006) Monitoring mesenchymal stromal cell developmental stage to apply on-time mechanical stimulation for ligament tissue engineering. *Tissue Eng* **12**: 3085-3095.
- Cho TJ, Kim J, Kwon SK, Oh K, Lee J, Lee DS, Cho J, Park SB (2012) A potent small-molecule inducer of chondrogenic differentiation of human bone marrow-derived mesenchymal stem cells. *Chem Sci* **3**: 3071-3075.
- Choi KM, Seo YK, Yoon HH, Song KY, Kwon SY, Lee HS, Park JK (2008) Effect of ascorbic acid on bone marrow-derived mesenchymal stem cell proliferation and differentiation. *J Biosci Bioeng* **105**: 586-594.
- Chowdhury F, Na S, Li D, Poh YC, Tanaka TS, Wang F, Wang N (2010) Material properties of the cell dictate stress-induced spreading and differentiation in embryonic stem cells. *Nat Mater* **9**: 82-88.
- Cigan AD, Nims RJ, Albro MB, Esau JD, Dreyer MP, Vunjak-Novakovic G, Hung CT, Ateshian GA, (2013) Insulin, ascorbate, and glucose have a much greater influence than transferrin and selenous acid on the *in vitro* growth of engineered cartilage in chondrogenic media. *Tissue Eng Part A* **19**: 1941-1948.
- Clark JM, Norman A, Notzli H (1997) Postnatal development of the collagen matrix in rabbit tibial plateau articular cartilage. *J Anat* **191**: 215-227.
- Coleman CM, Tuan RS (2003) Functional role of growth/differentiation factor 5 in chondrogenesis of limb mesenchymal cells. *Mech Dev* **120**: 823-836.
- Coleman CM, Vaughan EE, Browe DC, Mooney E, Howard L, Barry F (2013) Growth differentiation factor-5 enhances *in vitro* mesenchymal stromal cell chondrogenesis and hypertrophy. *Stem Cells Dev* **22**: 1968-1976.
- Derfoul A, Perkins GL, Hall DJ, Tuan RS (2006) Glucocorticoids promote chondrogenic differentiation of adult human mesenchymal stem cells by enhancing

expression of cartilage extracellular matrix genes. *Stem Cells* **24**: 1487-1495.

Dessau W, von der Mark H, von der Mark K, Fischer S (1980) Changes in the patterns of collagens and fibronectin during limb-bud chondrogenesis. *J Embryol Exp Morphol* **57**: 51-60.

Du J, Chen X, Liang X, Zhang G, Xu J, He L, Zhan Q, Feng XQ, Chien S, Yang C (2011) Integrin activation and internalization on soft ECM as a mechanism of induction of stem cell differentiation by ECM elasticity. *Proc Natl Acad Sci U S A* **108**: 9466-9471.

Engler AJ, Sen S, Sweeney HL, Discher DE (2006) Matrix elasticity directs stem cell lineage specification. *Cell* **126**: 677-689.

Enochson L, Stenberg J, Brittberg M, Lindahl A (2014) GDF5 reduces MMP13 expression in human chondrocytes *via* DKK1 mediated canonical Wnt signaling inhibition. *Osteoarthritis Cartilage* **22**: 566-577.

Enomoto-Iwamoto M, Iwamoto M, Mukudai Y, Kawakami Y, Nohno T, Higuchi Y, Takemoto S, Ohuchi H, Noji S, Kurisu K (1998) Bone morphogenetic protein signaling is required for maintenance of differentiated phenotype, control of proliferation, and hypertrophy in chondrocytes. *J Cell Biol* **140**: 409-418.

Estes BT, Gimble JM, Guilak F (2004) Mechanical signals as regulators of stem cell fate. *Curr Top Dev Biol* **60**: 91-126.

Feng G, Wan Y, Balian G, Laurencin CT, Li X (2008) Adenovirus-mediated expression of growth and differentiation factor-5 promotes chondrogenesis of adipose stem cells. *Growth Factors* **26**: 132-142.

Florine EM, Miller RE, Porter RM, Evans CH, Kurz B, Grodzinsky AJ (2013) Effects of dexamethasone on mesenchymal stromal cell chondrogenesis and aggrecanase activity: comparison of agarose and self-assembling peptide scaffolds *Cartilage* **4**: 63-74.

Franceschi RT (1992) The role of ascorbic acid in mesenchymal differentiation. *Nutr Rev* **50**: 65-70.

Furumatsu T, Ozaki T (2010) Epigenetic regulation in chondrogenesis. *Acta Med Okayama* **64**: 155-161.

Furumatsu T, Tsuda M, Taniguchi N, Tajima Y, Asahara H (2005) Smad3 induces chondrogenesis through the activation of SOX9 *via* CREB-binding protein/p300 recruitment. *J Biol Chem* **280**: 8343-8350.

Ge ZG, Hu Y, Heng BC, Yang Z, Ouyang HW, Lee EH, Cao T (2006) Osteoarthritis and therapy. *Arthritis Care Res* **55**: 493-500.

Geshi M, Takenouchi N, Yamauchi N, Nagai T (2000) Effects of sodium pyruvate in nonserum maturation medium on maturation, fertilization, and subsequent development of bovine oocytes with or without cumulus cells. *Biol Reprod* **63**: 1730-1734.

Goldring MB, Tsuchimochi K, Ijiri K (2006) The control of chondrogenesis. *J Cell Biochem* **97**: 33-44.

Gorlin JB, Yamin R, Egan S, Stewart M, Stossel TP, Kwiatkowski DJ, Hartwig JH (1990) Human endothelial actin-binding protein (ABP-280, nonmuscle filamin): a molecular leaf spring. *J Cell Biol* **111**: 1089-1105.

Han YS, Bang OS, Jin EJ, Park JH, Sonn JK, Kang SS (2004) High dose of glucose promotes chondrogenesis

via PKC α and MAPK signaling pathways in chick mesenchymal cells. *Cell Tissue Res* **318**: 571-578.

Handorf AM, Li WJ (2014) Induction of mesenchymal stem cell chondrogenesis through sequential administration of growth factors within specific temporal windows. *J Cell Physiol* **229**: 162-171.

Hassel S, Schmitt S, Hartung A, Roth M, Nohe A, Petersen N, Ehrlich M, Henis YI, Sebald W, Knaus P (2003) Initiation of Smad-dependent and Smad-independent signaling *via* distinct BMP-receptor complexes. *J Bone Joint Surg Am* **85-A Suppl 3**: 44-51.

Hatakeyama Y, Tuan RS, Shum L (2004) Distinct functions of BMP4 and GDF5 in the regulation of chondrogenesis. *J Cell Biochem* **91**: 1204-1217.

Hattori T, Ide H (1984) Limb bud chondrogenesis in cell culture, with particular reference to serum concentration in the culture medium. *Exp Cell Res* **150**: 338-346.

Healy C, Uwanogho D, Sharpe PT (1999) Regulation and role of Sox9 in cartilage formation. *Dev Dyn* **215**: 69-78.

Heine U, Munoz EF, Flanders KC, Ellingsworth LR, Lam HY, Thompson NL, Roberts AB, Sporn MB (1987) Role of transforming growth factor- β in the development of the mouse embryo. *J Cell Biol* **105**: 2861-2876.

Hellingman CA, Davidson EN, Koevoet W, Vitters EL, van den Berg WB, van Osch GJ, van der Kraan PM (2011) Smad signaling determines chondrogenic differentiation of bone-marrow-derived mesenchymal stem cells: inhibition of Smad1/5/8P prevents terminal differentiation and calcification. *Tissue Eng Part A* **17**: 1157-1167.

Hellingman CA, Koevoet W, Kops N, Farrell E, Jahr H, Liu W, Baatenburg de Jong RJ, Frenz DA, van Osch GJ (2010) Fibroblast growth factor receptors in *in vitro* and *in vivo* chondrogenesis: relating tissue engineering using adult mesenchymal stem cells to embryonic development. *Tissue Eng Part A* **16**: 545-556.

Hennig T, Lorenz H, Thiel A, Goetzke K, Dickhut A, Geiger F, Richter W (2007) Reduced chondrogenic potential of adipose tissue derived stromal cells correlates with an altered TGF β receptor and BMP profile and is overcome by BMP-6. *J Cell Physiol* **211**: 682-691.

Herlofsen S, Bryne J, Høiby T, Wang L, Issner R, Zhang X, Coyne M, Boyle P, Gu H, Meza-Zepeda L, Collas P, Mikkelsen T, Brinchmann J (2013) Genome-wide map of quantified epigenetic changes during *in vitro* chondrogenic differentiation of primary human mesenchymal stem cells. *BMC Genomics* **14**: 1-18.

Hogan BL (1996) Bone morphogenetic proteins: multifunctional regulators of vertebrate development. *Genes Dev* **10**: 1580-1594.

Holtzer H (1952) An experimental analysis of the development of the spinal column. Part I. response of pre-cartilage cells to size variations of the spinal cord. *J Exp Zool* **121**: 121-147.

Holtzer H (1964) Control of chondrogenesis in the embryo. *Biophys J* **4**: 239-250.

Holtzer H, Detwiler SR (1953) An experimental analysis of the development of the spinal column. III. induction of skeletogenous cells. *J Exp Zool* **123**: 335-369.

Hommes FA, Van Leeuwen G, Zilliken F (1962) Induction of cell differentiation: II. the isolation of a

chondrogenic factor from embryonic chick spinal cords and notochords. *Biochim Biophys Acta* **56**: 320-325.

Hong E, Reddi AH (2012) MicroRNAs in chondrogenesis, articular cartilage, and osteoarthritis: implications for tissue engineering. *Tissue Eng Part B Rev* **18**: 445-453.

Houlihan DD, Mabuchi Y, Morikawa S, Niibe K, Araki D, Suzuki S, Okano H, Matsuzaki Y (2012) Isolation of mouse mesenchymal stem cells on the basis of expression of sca-1 and pdgfr- α . *Nat Protoc* **7**: 2103-2111.

Huang AH, Farrell MJ, Mauck RL (2010a) Mechanics and mechanobiology of mesenchymal stem cell-based engineered cartilage. *J Biomech* **43**: 128-136.

Huang AH, Stein A, Mauck RL (2010b) Evaluation of the complex transcriptional topography of mesenchymal stem cell chondrogenesis for cartilage tissue Engineering. *Tissue Eng Part A* **16**: 2699-2708.

Huey DJ, Hu JC, Athanasiou KA (2012) Unlike bone, cartilage regeneration remains elusive. *Science* **338**: 917-921.

Hunziker EB (2009) The elusive path to cartilage regeneration. *Adv Mater* **21**: 3419-3424.

Hunziker EB, Kapfinger E, Geiss J (2007) The structural architecture of adult mammalian articular cartilage evolves by a synchronized process of tissue resorption and neoformation during postnatal development. *Osteoarthritis Cartilage* **15**: 403-413.

Huse M, Chen Y-G, Massagué J, Kuriyan J (1999) Crystal structure of the cytoplasmic domain of the Type I TGF β receptor in complex with FKBP12. *Cell* **96**: 425-436.

Hyde G, Dover S, Aszodi A, Wallis GA, Boot-Handford RP (2007) Lineage tracing using matrilin-1 gene expression reveals that articular chondrocytes exist as the joint interzone forms. *Dev Biol* **304**: 825-833.

Ito MM, Kida MY (2000) Morphological and biochemical re-evaluation of the process of cavitation in the rat knee joint: cellular and cell strata alterations in the interzone. *J Anat* **197**: 659-679.

Iwamoto M, Ohta Y, Larmour C, Enomoto-Iwamoto M (2013) Toward regeneration of articular cartilage. *Birth Defects Res C Embryo Today* **99**: 192-202.

Jakobsen RB, Østrup E, Zhang X, S.Mikkelsen T, E.Brinchmann J (2014) Analysis of the effects of five factors relevant to *in vitro* chondrogenesis of human mesenchymal stem cells using factorial design and high throughput mRNA-profiling. *PLoS ONE* **9**: art.no.e96615.

Jin E-J, Lee S-Y, Choi Y-A, Jung J-C, Bang O-S, Kang S-S (2006) BMP-2-enhanced chondrogenesis involves p38 MAPK-mediated down-regulation of Wnt-7a pathway. *Mol Cells* **22**: 353-359.

Johnson K, Zhu S, Tremblay MS, Payette JN, Wang J, Bouchez LC, Meeusen S, Althage A, Cho CY, Wu X, Schultz PG (2012) A stem cell-based approach to cartilage repair. *Science* **336**: 717-721.

Johnstone B, Hering TM, Caplan AI, Goldberg VM, Yoo JU (1998) *In vitro* chondrogenesis of bone marrow-derived mesenchymal progenitor cells. *Exp Cell Res* **238**: 265-272.

Kafienah W, Mistry S, Perry MJ, Politopoulou G, Hollander AP (2007) Pharmacological regulation of adult

stem cells: chondrogenesis can be induced using a synthetic inhibitor of the retinoic acid receptor. *Stem Cells* **25**: 2460-2468.

Kanichai M, Ferguson D, Prendergast PJ, Campbell VA (2008) Hypoxia promotes chondrogenesis in rat mesenchymal stem cells: a role for AKT and hypoxia-inducible factor (HIF)-1 α . *J Cell Physiol* **216**: 708-715.

Kassis I, Zangi L, Rivkin R, Levinsky, Samuel S, Marx G, Gorodetsky R (2006) Isolation of mesenchymal stem cells from G-CSF-mobilized human peripheral blood using fibrin microbeads. *Bone Marrow Transplant* **37**: 967-976.

Kato M, Takenawa T, Twardzik DR (1988) Effect of transforming growth factor- α on inositol phospholipid-metabolism in human epidermoid carcinoma-cells. *J Cell Biochem* **37**: 339-345.

Kim YJ, Kim HJ, Im GI (2008) PTHrP promotes chondrogenesis and suppresses hypertrophy from both bone marrow-derived and adipose tissue-derived MSCs. *Biochem Biophys Res Commun* **373**: 104-108.

Kock L, van Donkelaar CC, Ito K (2012) Tissue engineering of functional articular cartilage: the current status. *Cell Tissue Res* **347**: 613-627.

Kono T, Robinson FW, Blevins TL, Ezaki O (1982) Evidence that translocation of the glucose transport activity is the major mechanism of insulin action on glucose transport in fat cells. *J Biol Chem* **257**: 10942-10947.

Koyama E, Shibukawa Y, Nagayama M, Sugito H, Young B, Yuasa T, Okabe T, Ochiai T, Kamiya N, Rountree RB, Kingsley DM, Iwamoto M, Enomoto-Iwamoto M, Pacifici M (2008) A distinct cohort of progenitor cells participates in synovial joint and articular cartilage formation during mouse limb skeletogenesis. *Dev Biol* **316**: 62-73.

Kramer J, Hegert C, Guan K, Wobus AM, Müller PK, Rohwedel J (2000) Embryonic stem cell-derived chondrogenic differentiation *in vitro*: activation by BMP-2 and BMP-4. *Mech Dev* **92**: 193-205.

Kulyk WM, Upholt WB, Kosher RA (1989) Fibronectin gene expression during limb cartilage differentiation. *Development* **106**: 449-455.

Kurth T, Hedbom E, Shintani N, Sugimoto M, Chen FH, Haspl M, Martinovic S, Hunziker EB (2007) Chondrogenic potential of human synovial mesenchymal stem cells in alginate. *Osteoarthritis Cartilage* **15**: 1178-1189.

Lash JW, Holtzer H, Whitehouse MW (1960) *In vitro* studies on chondrogenesis; the uptake of radioactive sulfate during cartilage induction. *Dev Biol* **2**: 76-89.

Lash JW, Hommes FA, Zilliken F (1962) Induction of cell differentiation: I. The *in vitro* induction of vertebral cartilage with a low-molecular-weight tissue component. *Biochim Biophys Acta* **56**: 313-319.

Lee CH, Cook JL, Mendelson A, Moiola EK, Yao H, Mao JJ (2010) Regeneration of the articular surface of the rabbit synovial joint by cell homing: a proof of concept study. *Lancet* **376**: 440-448.

Lee HH, Chang CC, Shieh MJ, Wang JP, Chen YT, Young TH, Hung SC (2013) Hypoxia enhances chondrogenesis and prevents terminal differentiation through PI3K/Akt/FoxO dependent anti-apoptotic effect. *Sci Rep* **3**: 1-12.

- Lee JM, Im GI (2012) PTHrP isoforms have differing effect on chondrogenic differentiation and hypertrophy of mesenchymal stem cells. *Biochem Biophys Res Commun* **421**: 819-824.
- Lefebvre V, Li P, de Crombrughe B (1998) A new long form of Sox5 (L-Sox5), Sox6 and Sox9 are coexpressed in chondrogenesis and cooperatively activate the type II collagen gene. *EMBO J* **17**: 5718-5733.
- Lefebvre V, Smits P (2005) Transcriptional control of chondrocyte fate and differentiation. *Birth Defects Res C Embryo Today* **75**: 200-212.
- Leijten J, Georgi N, Teixeira LM, van Blitterswijk CA, Post JN, Karperien M (2014) Metabolic programming of mesenchymal stromal cells by oxygen tension directs chondrogenic cell fate. *Proc Natl Acad Sci USA* **111**: 13954-13959.
- Leijten JC, Emons J, Sticht C, van Gool S, Decker E, Uitterlinden A, Rappold G, Hofman A, Rivadeneira F, Scherjon S, Wit JM, van Meurs J, van Blitterswijk CA, Karperien M (2012) Gremlin 1, frizzled-related protein, and Dkk-1 are key regulators of human articular cartilage homeostasis. *Arthritis Rheum* **64**: 3302-3312.
- Li J, Ohliger J, Pei M (2014) Significance of epigenetic landscape in cartilage regeneration from the cartilage development and pathology perspective. *Stem Cells Dev* **23**: 1178-1194.
- Li J, Pei M (2012) Cell senescence: a challenge in cartilage engineering and regeneration. *Tissue Eng Part B Rev* **18**: 270-287.
- Li J, Zhao Z, Liu J, Huang N, Long D, Wang J, Li X, Liu Y (2010) MEK/ERK and p38 MAPK regulate chondrogenesis of rat bone marrow mesenchymal stem cells through delicate interaction with TGF- β 1/Smads pathway. *Cell Prolif* **43**: 333-343.
- Loeser RF (2009) Aging and osteoarthritis: the role of chondrocyte senescence and aging changes in the cartilage matrix. *Osteoarthritis Cartilage* **17**: 971-979.
- Lutolf MP, Gilbert PM, Blau HM (2009) Designing materials to direct stem-cell fate. *Nature* **462**: 433-441.
- Lutz M, Knaus P (2002) Integration of the TGF-beta pathway into the cellular signalling network. *Cell Signal* **14**: 977-988.
- Lysdahl H, Baatrup A, Nielsen A, Foldager C, Bunger C (2013) Phenol red inhibits chondrogenic differentiation and affects osteogenic differentiation of human mesenchymal stem cells *in vitro*. *Stem Cell Rev* **9**: 132-139.
- Mackay AM, Beck SC, Murphy JM, Barry FP, Chichester CO, Pittenger MF (1998) Chondrogenic differentiation of cultured human mesenchymal stem cells from marrow. *Tissue Eng* **4**: 415-428.
- Mackie EJ, Ahmed YA, Tatarczuch L, Chen KS, Mirams M (2008) Endochondral ossification: How cartilage is converted into bone in the developing skeleton. *Int J Biochem Cell Biol* **40**: 46-62.
- Maor G, Silbermann M, von der Mark K, Heingard D, Laron Z (1993) Insulin enhances the growth of cartilage in organ and tissue cultures of mouse neonatal mandibular condyle. *Calcif Tissue Int* **52**: 291-299.
- Massagu J (1998) TGF- β signal transduction. *Annu Rev Biochem* **67**: 753-791.
- Massagu J, Wotton D (2000) Transcriptional control by the TGF- β /Smad signaling system. *EMBO J* **19**: 1745-1754.
- Mastrogiacomo M, Cancedda R, Quarto R (2001) Effect of different growth factors on the chondrogenic potential of human bone marrow stromal cells. *Osteoarthritis Cartilage* **9**: S36-S40.
- Mau E, Whetstone H, Yu C, Hopyan S, Wunder JS, Alman BA (2007) PTHrP regulates growth plate chondrocyte differentiation and proliferation in a Gli3 dependent manner utilizing hedgehog ligand dependent and independent mechanisms. *Dev Biol* **305**: 28-39.
- Meretoja VV, Dahlin RL, Kasper FK, Mikos AG (2012) Enhanced chondrogenesis in co-cultures with articular chondrocytes and mesenchymal stem cells. *Biomaterials* **33**: 6362-6369.
- Minowada G, Jarvis LA, Chi CL, Neubuser A, Sun X, Hacoen N, Krasnow MA, Martin GR (1999) Vertebrate Sprouty genes are induced by FGF signaling and can cause chondrodysplasia when overexpressed. *Development* **126**: 4465-4475.
- Morikawa S, Mabuchi Y, Niibe K, Suzuki S, Nagoshi N, Sunabori T, Shimmura S, Nagai Y, Nakagawa T, Okano H, Matsuzaki Y (2009) Development of mesenchymal stem cells partially originate from the neural crest. *Biochem Biophys Res Commun* **379**: 1114-1119.
- Mueller M, Blunk T, Appel B, Maschke A, Goepferich A, Zellner J, Englert C, Prantl L, Kujat R, Nerlich M, Angele P (2013a) Insulin is essential for *in vitro* chondrogenesis of mesenchymal progenitor cells and influences chondrogenesis in a dose-dependent manner. *Int Orthop* **37**: 153-158.
- Mueller MB, Fischer M, Zellner J, Berner A, Dienstknecht T, Kujat R, Prantl L, Nerlich M, Tuan RS, Angele P (2013b) Effect of parathyroid hormone-related protein in an *in vitro* hypertrophy model for mesenchymal stem cell chondrogenesis. *Int Orthop* **37**: 945-951.
- Mueller MB, Tuan RS (2008) Functional characterization of hypertrophy in chondrogenesis of human mesenchymal stem cells. *Arthritis Rheum* **58**: 1377-1388.
- Murphy JM, Dixon K, Beck S, Fabian D, Feldman A, Barry F (2002) Reduced chondrogenic and adipogenic activity of mesenchymal stem cells from patients with advanced osteoarthritis. *Arthritis Rheum* **46**: 704-713.
- Murphy MK, Huey DJ, Hu JC, Athanasiou KA (2015) TGF- β 1, GDF-5, and BMP-2 stimulation induces chondrogenesis in expanded human articular chondrocytes and marrow-derived stromal cells. *Stem Cells* **33**: 762-773.
- Murtaugh LC, Chyung JH, Lassar AB (1999) Sonic hedgehog promotes somitic chondrogenesis by altering the cellular response to BMP signaling. *Genes Dev* **13**: 225-237.
- Ned RM, Swat W, Andrews NC (2003) Transferrin receptor 1 is differentially required in lymphocyte development. *Blood* **102**: 3711-3718.
- Ng F, Boucher S, Koh S, Sastry KS, Chase L, Lakshminpathy U, Choong C, Yang Z, Vemuri MC, Rao MS (2008) PDGF, TGF- β , and FGF signaling is important for differentiation and growth of mesenchymal stem cells (MSCs): transcriptional profiling can identify markers and

signaling pathways important in differentiation of MSCs into adipogenic, chondrogenic, and osteogenic lineages. *Blood* **112**: 295-307.

Nonaka K, Shum L, Takahashi I, Takahashi K, Ikura T, Dashner R, Nuckolls GH, Slavkin HC (1999) Convergence of the BMP and EGF signaling pathways on Smad1 in the regulation of chondrogenesis. *Int J Dev Biol* **43**: 795-807.

O'Connor CJ, Case N, Guilak F (2013) Mechanical regulation of chondrogenesis. *Stem Cell Res Ther* **4**: 61.

O'Donnell-Tormey J, Nathan CF, Lanks K, DeBoer CJ, de la Harpe J (1987) Secretion of pyruvate. An antioxidant defense of mammalian cells. *J Exp Med* **165**: 500-514.

Oakley RH, Jewell CM, Yudit MR, Bofetiado DM, Cidlowski JA (1999) The dominant negative activity of the human glucocorticoid receptor beta isoform. *J Biol Chem* **274**: 27857-27866.

Oldershaw RA, Baxter MA, Lowe ET, Bates N, Grady LM, Soncin F, Brison DR, Hardingham TE, Kimber SJ (2010) Directed differentiation of human embryonic stem cells toward chondrocytes. *Nat Biotechnol* **28**: 1187-1194.

Olsen BR, Reginato AM, Wang W (2000) Bone development. *Annu Rev Cell Dev Biol* **16**: 191-220.

Onyekwelu I, Goldring MB, Hidaka C (2009) Chondrogenesis, joint formation, and articular cartilage regeneration. *J Cell Biochem* **107**: 383-392.

Oszycza AM, Diefenderfer DL, Bhargava G, Leboy PS (2004) Different effects of BMP-2 on marrow stromal cells from human and rat bone. *Cells Tissues Organs* **176**: 109-119.

Otte P (1991) Basic cell metabolism of articular cartilage. Manometric studies. *Z Rheumatol* **50**: 304-312.

Owens EM, Solursh M (1982) Cell-cell interaction by mouse limb cells during *in vitro* chondrogenesis: Analysis of the brachypod mutation. *Dev Biol* **91**: 376-388.

Oze H, Hirao M, Ebina K, Shi K, Kawato Y, Kaneshiro S, Yoshikawa H, Hashimoto J (2012) Impact of medium volume and oxygen concentration in the incubator on pericellular oxygen concentration and differentiation of murine chondrogenic cell culture. *In Vitro Cell Dev Biol Anim* **48**: 123-130.

Pacifici M, Koyama E, Iwamoto M (2005) Mechanisms of synovial joint and articular cartilage formation: Recent advances, but many lingering mysteries. *Birth Defects Res* **75**: 237-248.

Padh H (1991) Vitamin C: newer insights into its biochemical functions. *Nutr Rev* **49**: 65-70.

Paik S, Jung HS, Lee S, Yoon DS, Park MS, Lee JW (2012) miR-449a regulates the chondrogenesis of human mesenchymal stem cells through direct targeting of lymphoid enhancer-binding factor-1. *Stem Cells Dev* **21**: 3298-3308.

Park JS, Chu JS, Tsou AD, Diop R, Tang Z, Wang A, Li S (2011) The effect of matrix stiffness on the differentiation of mesenchymal stem cells in response to TGF- β . *Biomaterials* **32**: 3921-3930.

Park Y, Sugimoto M, Watrin A, Chiquet M, Hunziker EB (2005) BMP-2 induces the expression of chondrocyte-specific genes in bovine synovium-derived progenitor cells cultured in three-dimensional alginate hydrogel. *Osteoarthritis Cartilage* **13**: 527-536.

Pelttari K, Winter A, Steck E, Goetzke K, Hennig T, Ochs BG, Aigner T, Richter W (2006) Premature induction of hypertrophy during *in vitro* chondrogenesis of human mesenchymal stem cells correlates with calcification and vascular invasion after ectopic transplantation in SCID mice. *Arthritis Rheum* **54**: 3254-3266.

Peterkofsky B (1972) The effect of ascorbic acid on collagen polypeptide synthesis and proline hydroxylation during the growth of cultured fibroblasts. *Arch Biochem Biophys* **152**: 318-328.

Peterkofsky B (1991) Ascorbate requirement for hydroxylation and secretion of procollagen: relationship to inhibition of collagen synthesis in scurvy. *Am J Clin Nutr* **54**: 1135S-1140S.

Pizette S, Niswander L (2000) BMPs are required at two steps of limb chondrogenesis: formation of prechondrogenic condensations and their differentiation into chondrocytes. *Dev Biol* **219**: 237-249.

Potdar PD, D'Souza SB (2010) Ascorbic acid induces *in vitro* proliferation of human subcutaneous adipose tissue derived mesenchymal stem cells with upregulation of embryonic stem cell pluripotency markers Oct4 and SOX 2. *Hum Cell* **23**: 152-155.

Quarto R, Campanile G, Cancedda R, Dozin B (1992) Thyroid hormone, insulin, and glucocorticoids are sufficient to support chondrocyte differentiation to hypertrophy: a serum-free analysis. *J Cell Biol* **119**: 989-995.

Rabie AB, Tang GH, Xiong H, Hagg U (2003) PTHrP regulates chondrocyte maturation in condylar cartilage. *J Dent Res* **82**: 627-631.

Rainbow RS, Kwon H, Foote AT, Preda RC, Kaplan DL, Zeng L (2013) Muscle cell-derived factors inhibit inflammatory stimuli-induced damage in hMSC-derived chondrocytes. *Osteoarthritis Cartilage* **21**: 990-998.

Randua TM, Schildberg FA, Alini M, Wimmer MD, Haddouti EM, Gravius S, Ito K, Stoddart MJ (2013) The effect of dexamethasone and triiodothyronine on terminal differentiation of primary bovine chondrocytes and chondrogenically differentiated mesenchymal stem cells. *PLoS One* **8**.

Retting KN, Song B, Yoon BS, Lyons KM (2009) BMP canonical Smad signaling through Smad1 and Smad5 is required for endochondral bone formation. *Development* **136**: 1093-1104.

Rosen O (1987) After insulin binds. *Science* **237**: 1452-1458.

Roth SY, Denu JM, Allis CD (2001) Histone acetyltransferases. *Annu Rev Biochem* **70**: 81-120.

Sanders EJ (1988) The roles of epithelial-mesenchymal cell interactions in developmental processes. *Biochem Cell Biol* **66**: 530-540.

Schäfer R, Kehlbach R, Wiskirchen J, Bantleon Rd, Pintaske Jr, Brehm BR, Gerber A, Wolburg H, Claussen CD, Northoff H (2007) Transferrin receptor upregulation: *in vitro* labeling of rat mesenchymal stem cells with superparamagnetic iron oxide 1. *Radiology* **244**: 514-523.

Schmierer B, Hill CS (2007) TGF β -SMAD signal transduction: molecular specificity and functional flexibility. *Nat Rev Mol Cell Biol* **8**: 970-982.

- Schugar R, Robbins P, Deasy B (2008) Small molecules in stem cell self-renewal and differentiation. *Gene Ther* **15**: 126-135.
- Sekiya I, Larson BL, Vuoristo JT, Reger RL, Prockop DJ (2005) Comparison of effect of BMP-2,-4, and-6 on *in vitro* cartilage formation of human adult stem cells from bone marrow stroma. *Cell Tissue Res* **320**: 269-276.
- Serra R, Johnson M, Filvaroff EH, LaBorde J, Sheehan DM, Derynck R, Moses HL (1997) Expression of a truncated, kinase-defective TGF-beta type II receptor in mouse skeletal tissue promotes terminal chondrocyte differentiation and osteoarthritis. *J Cell Biol* **139**: 541-552.
- Seyedin SM, Thompson AY, Bentz H, Rosen DM, McPherson JM, Conti A, Siegel NR, Galluppi GR, Piez KA (1986) Cartilage-inducing factor-A. Apparent identity to transforming growth factor-beta. *J Biol Chem* **261**: 5693-5695.
- Shakibaei M (1998) Inhibition of chondrogenesis by Integrin antibody *in vitro*. *Exp Cell Res* **240**: 95-106.
- Shen B, Wei A, Whittaker S, Williams LA, Tao H, Ma DD, Diwan AD (2010) The role of BMP-7 in chondrogenic and osteogenic differentiation of human bone marrow multipotent mesenchymal stromal cells *in vitro*. *J Cell Biochem* **109**: 406-416.
- Shikhman AR, Brinson DC, Valbracht J, Lotz MK (2001) Cytokine regulation of facilitated glucose transport in human articular chondrocytes. *J Immunol* **167**: 7001-7008.
- Shintani N, Hunziker EB (2011) Differential effects of dexamethasone on the chondrogenesis of mesenchymal stromal cells: influence of microenvironment, tissue origin and growth factor. *Eur Cell Mater* **22**: 302-319; discussion 319-320.
- Shintani N, Kurth T, Hunziker EB (2007) Expression of cartilage-related genes in bovine synovial tissue. *J Orthop Res* **25**: 813-819.
- Shu J, Deng H (2013) Lineage specifiers: new players in the induction of pluripotency. *Genomics Proteomics Bioinformatics* **11**: 259-263.
- Shukunami C, Shigeno C, Atsumi T, Ishizeki K, Suzuki F, Hiraki Y (1996) Chondrogenic differentiation of clonal mouse embryonic cell line ATDC5 *in vitro*: differentiation-dependent gene expression of parathyroid hormone (PTH)/PTH-related peptide receptor. *J Cell Biol* **133**: 457-468.
- Solchaga LA, Penick K, Porter JD, Goldberg VM, Caplan AI, Welter JF (2005) FGF-2 enhances the mitotic and chondrogenic potentials of human adult bone marrow-derived mesenchymal stem cells. *J Cell Physiol* **203**: 398-409.
- Spencer JA, Ferraro F, Roussakis E, Klein A, Wu J, Runnels JM, Zaher W, Mortensen LJ, Alt C, Turcotte R, Yusuf R, Cote D, Vinogradov SA, Scadden DT, Lin CP (2014) Direct measurement of local oxygen concentration in the bone marrow of live animals. *Nature* **508**: 269-273.
- Squillaro T, Severino V, Alessio N, Farina A, Di Bernardo G, Cipollaro M, Peluso G, Chambery A, Galderisi U (2015) De-regulated expression of the BRG1 chromatin remodeling factor in bone marrow mesenchymal stromal cells induces senescence associated with the silencing of NANOG and changes in the levels of chromatin proteins. *Cell Cycle* **14**: 1315-1326.
- Stone N, Meister A (1962) Function of ascorbic acid in the conversion of proline to collagen hydroxyproline. *Nature* **194**: 555-557.
- Storm EE, Huynh TV, Copeland NG, Jenkins NA, Kingsley DM, Lee S-J (1994) Limb alterations in brachypodism mice due to mutations in a new member of the TGF[beta]-superfamily. *Nature* **368**: 639-643.
- Studer D, Millan C, Oeztuerk E, Maniura-Weber K, Zenobi-Wong M (2012) Molecular and biophysical mechanisms regulating hypertrophic differentiation in chondrocytes and mesenchymal stem cells. *Eur Cell Mater* **24**: 118-135.
- Sun Z, Zhang Y, Yang S, Jia J, Ye S, Chen D, Mo F (2012) Growth differentiation factor 5 modulation of chondrogenesis of self-assembled constructs involves gap junction-mediated intercellular communication. *Dev Growth Differ* **54**: 809-817.
- Takashima Y, Era T, Nakao K, Kondo S, Kasuga M, Smith AG, Nishikawa S (2007) Neuroepithelial cells supply an initial transient wave of MSC differentiation. *Cell* **129**: 1377-1388.
- Tate MLK, Falls TD, McBride SH, Atit R, Knothe UR (2008) Mechanical modulation of osteochondroprogenitor cell fate. *Int J Biochem Cell Biol* **40**: 2720-2738.
- Tsai TL, Manner PA, Li WJ (2013) Regulation of mesenchymal stem cell chondrogenesis by glucose through protein kinase C/transforming growth factor signaling. *Osteoarthritis Cartilage* **21**: 368-376.
- Tsutsumi S, Shimazu A, Miyazaki K, Pan H, Koike C, Yoshida E, Takagishi K, Kato Y (2001) Retention of multilineage differentiation potential of mesenchymal cells during proliferation in response to FGF. *Biochem Biophys Res Commun* **288**: 413-419.
- Tuli R, Tuli S, Nandi S, Huang X, Manner PA, Hozack WJ, Danielson KG, Hall DJ, Tuan RS (2003) Transforming growth factor-beta-mediated chondrogenesis of human mesenchymal progenitor cells involves N-cadherin and mitogen-activated protein kinase and Wnt signaling cross-talk. *J Biol Chem* **278**: 41227-41236.
- Vacanti V, Kong E, Suzuki G, Sato K, Cauty JM, Lee TC (2005) Phenotypic changes of adult porcine mesenchymal stem cells induced by prolonged passaging in culture. *J Cell Physiol* **205**: 194-201.
- Vater C, Kasten P, Stiehler M (2011) Culture media for the differentiation of mesenchymal stromal cells. *Acta Biomater* **7**: 463-477.
- Vortkamp A, Lee K, Lanske B, Segre GV, Kronenberg HM, Tabin CJ. (1996) Regulation of rate of cartilage differentiation by Indian hedgehog and PTH-related protein. *Science* **273**: 613-622.
- Wang CY, Chen LL, Kuo PY, Chang JL, Wang YJ, Hung SC (2010) Apoptosis in chondrogenesis of human mesenchymal stem cells: effect of serum and medium supplements. *Apoptosis* **15**: 439-449.
- Wang JA, Chen TL, Jiang J, Shi H, Gui C, Luo RH, Xie XJ, Xiang MX, Zhang X (2008) Hypoxic preconditioning attenuates hypoxia/reoxygenation-induced apoptosis in mesenchymal stem cells. *Acta Pharmacol Sin* **29**: 74-82.
- Washington JM, Rathjen J, Felquer F, Lonic A, Bettess MD, Hamra N, Semendric L, Tan BS, Lake JA, Keough RA, Morris MB, Rathjen PD (2010) L-Proline induces

differentiation of ES cells: a novel role for an amino acid in the regulation of pluripotent cells in culture. *Am J Physiol Cell Physiol* **298**: 982-992.

Wei X, Wright GC, Sokoloff L (1986) The effect of sodium selenite on chondrocytes in monolayer culture. *Arthritis Rheum* **29**: 660-664.

Weiss S, Hennig T, Bock R, Steck E, Richter W (2010) Impact of growth factors and PTHrP on early and late chondrogenic differentiation of human mesenchymal stem cells. *J Cell Physiol* **223**: 84-93.

Wilson L, Yang Q, Szustakowski JD, Gullicksen PS, Halse R (2007) Pyruvate induces mitochondrial biogenesis by a PGC-1 α -independent mechanism. *Am J Physiol Cell Physiol* **292**: 1599-1605.

Wise JK, Yarin AL, Megaridis CM, Cho M (2009) Chondrogenic differentiation of human mesenchymal stem cells on oriented nanofibrous scaffolds: engineering the superficial zone of articular cartilage. *Tissue Eng Part A* **15**: 913-921.

Wu L, Prins HJ, Helder MN, van Blitterswijk CA, Karperien M (2012) Trophic effects of mesenchymal stem cells in chondrocyte co-cultures are independent of culture conditions and cell sources. *Tissue Eng Part A* **18**: 1542-1551.

Wu YN, Law JBK, He AY, Low HY, Hui JHP, Lim CT, Yang Z, Lee EH (2014) Substrate topography determines the fate of chondrogenesis from human mesenchymal stem cells resulting in specific cartilage phenotype formation. *Nanomedicine* **10**: 1507-1516.

Yan J, Tian J, Zheng Y, Han Y, Lu S (2012) Selenium promotes proliferation of chondrogenic cell ATDC5

by increment of intracellular ATP content under serum deprivation. *Cell Biochem Funct* **30**: 657-663.

Yang X, Trehan SK, Guan Y, Sun C, Moore DC, Jayasuriya CT, Chen Q (2014) Matrilin-3 inhibits chondrocyte hypertrophy as a bone morphogenetic protein-2 antagonist. *J Biol Chem* **289**: 34768-34779.

Yoshida Y, Yamanaka S (2010) Recent stem cell advances: induced pluripotent stem cells for disease modeling and stem cell-based regeneration. *Circulation* **122**: 80-87.

Zerega B, Cermelli S, Bianco P, Cancedda R, Cancedda FD (1999) Parathyroid Hormone[PTH(1-34)] and parathyroid hormone-related protein[PTHrP(1-34)] promote reversion of hypertrophic chondrocytes to a prehypertrophic proliferating phenotype and prevent terminal differentiation of osteoblast-like cells. *J Bone Miner Res* **14**: 1281-1289.

Zhang J, Wu Y, Thote T, Lee EH, Ge Z, Yang Z (2014) The influence of scaffold microstructure on chondrogenic differentiation of mesenchymal stem cells. *Biomed Mater* **9**: 035011.

Zhang Y, Ryan JA, Di Cesare PE, Liu J, Walsh CA, You Z (2007) Doublecortin is expressed in articular chondrocytes. *Biochem Biophys Res Commun* **363**: 694-700.

Editor's Note: All questions/comments by the reviewers were answered by text changes. Therefore, there is no Discussion with Reviewers section.

Scientific Editor in charge of the paper: Mauro Alini.