



DIVERSE EFFECTS OF PULSED ELECTRICAL STIMULATION ON CELLS – WITH A FOCUS ON CHONDROCYTE AND CARTILAGE REGENERATION

T. Ning^{1,2,5}, K. Zhang^{1,3}, B. Chin Heng⁴ and Z. Ge^{1,*}

¹Department of Biomedical Engineering, College of Engineering, Peking University, Beijing 100871, China
²Peking-Tsinghua Center for Life Sciences, Beijing, China
³Chinese Academy of Sciences Investment Management Co., Ltd., Beijing, 100080, China
⁴School of Stomatology, Peking University, Beijing 100081, Beijing, China
⁵Academy for Advanced Interdisciplinary Studies, Peking University, Beijing, 100871, China

Abstract

Biological effects of pulsed electrical stimulation (PES) on cells and tissues have been intensively studied with the aim of advancing their biomedical applications. These effects vary significantly depending on PES parameters, cell and tissue types, which can be attributed to the diverse variety of signaling pathways, ion channels, and epigenetic mechanisms involved. The development of new technology platforms, such as nanosecond pulsed electric fields (nsPEFs) with finely tuned parameters, have added further complexity. The present review systematically examines current research progress in various aspects of PES, from physical models to biological effects on cells and tissues, including voltage-sensing domains of voltage-gated channels, pore formation, intracellular components/organelles, and signaling pathways. Emphasis is placed on the complexity of PES parameters and inconsistency of induced biological effects, with the aim of exploring the underlying physical and cellular mechanisms of the physiological effects of electrical stimulation on cells. With chondrogenic differentiation of stem cells and cartilage regeneration as examples, the underlying mechanisms involved were reviewed and analyzed, hoping to move forward towards potential biomedical applications. Hopefully, the present review will inspire more interest in the wider clinical applications of PES and lay the basis for further comprehensive studies in this field.

Keywords: Electrical stimulation, electric fields, chondrocytes, chondrogenic differentiation, cartilage, stem cells.

*Address for correspondence: Dr Zigang Ge, Department of Biomedical Engineering, College of Engineering, Peking University, Beijing, 100871, China.

Telephone number: +86 01062756736 Fax number: +86 01062751812 Email: gez@pku.edu.cn

Copyright policy: This article is distributed in accordance with Creative Commons Attribution Licence (http://creativecommons.org/licenses/by-sa/4.0/).

	List of Abbreviations	Cx43 DNMT1	connexin 43 DNA methyltransferase 1
ADSCs	adipose-derived stem cells	ERK	extracellular-signal-regulated kinase
AGG	aggrecan	FDA	Food and Drug Administration
Bax	Bcl-2-associated X protein	FGF2	fibroblast growth factor 2
Bcl-2	B-cell lymphoma 2	GAG	glycosaminoglycan
BMI	body mass index	h-TENS	high-frequency transcutaneous
BMPs	bone morphogenetic proteins		electrical nerve stimulation
BMSCs	bone marrow stromal cells	IFC	interferential current
CaM	calmodulin	IGF2	insulin-like growth factor II
CaN-NFAT	calcineurin-nuclear factor of	IL	interleukin
	activated T cell	IRS	insulin receptor substrate
cGMP	cyclic guanosine monophosphate	1-TENS	low-frequency transcutaneous
COL II	collagen II		electrical nerve stimulation
CVI	chronic venous insufficiency	JNK	c-Jun N-terminal kinase

МАРК	mitogen-activated kinase
MSCs	mesenchymal stem cells
NEUROG2	neurogenin 2
NMES	neuromuscular electrical stimulation
NIN	non-invasive interactive
	neurostimulation
NO	nitric oxide
nsPEFs	nanosecond-pulsed electric fields
OA	osteoarthritis
PD-ECGF	platelet-derived endothelial cell
	growth factor
PEG	polyethylene glycol
PES	pulsed electrical stimulation
PIP2	phosphatidylinositol
	4,5-bisphosphate
ROS	reactive oxygen species
snRNPs	small nuclear ribonucleoprotein
	particles
TGF-β	transforming growth factor beta 1
VEGF	vascular endothelial growth factor
VOCCs	voltage-operated calcium channels
VSDs	voltage-sensing domains
	0 0

Introduction

PES has long been studied in the context of clinical therapy for various diseases, including wound healing (Ud-Din and Bayat, 2014), bone and cartilage regeneration (Chao and Inoue, 2003; Ciombor and Aaron, 2005), recovery of motor neurons (Rossini et al., 2015), contractile properties of infarcted heart tissues (Hirt et al., 2014), and rehabilitation of muscle contractile function (Kern et al., 2016), as well as tumor therapy through triggering permanent membrane lysis or loss of homeostasis (Cemazar et al., 2013). Several PES devices have been approved by the US FDA for use in the treatment of various conditions, such as tumors, arthritis, and pain, with durations varying from μ s to ms (Table 1). Accumulation of Joule heating energy is still a major hurdle for clinical application of electrical stimulation (Voldman, 2006; Weaver et al., 1999), which in turn demands critical restrictions on PES parameters, even though mammalian cells can survive up to a 1-3 °C increase in temperature. Relatively broad or unfocused biological targets limit the exploration of the mechanistic effects of electrical stimulation, as well as their widespread applications in clinical therapy (Okino and Mohri, 1987; Yao et al., 2009). Some reviews have also focused on treatment of conditions such as OA, fracture healing, and muscle pain by PES (Ebrahim et al., 2014; Negm et al., 2013). PES has also been used for plant and animal tissue drying processes and food preservation (Toepfl et al., 2006). Furthermore, high intensity PES can be used as a highly effective process for decontamination of liquid food (Sobrino-López and Martín-Belloso, 2010).

The application of PES, with varying parameters, may lead to significantly varied outcomes under

different physiological conditions (depending on cell phenotypes, attachment, and extracellular environment), which limits wider biomedical useage. For example, PES with voltages ≥ 3 V and frequencies ≥ 130 Hz has been reported to ameliorate Parkinson's disease, while PES with a frequency of 5 Hz significantly worsens akinesia (Moro *et al.*, 2002). PES has been reported to prevent atrophy of long-term denervated muscles, resulting in increased muscle cross-sectional area (Ashley *et al.*, 2007). On the other hand, PES (frequency of 20 Hz and durations longer than 2 ms) can also lead to atrophy of muscle fibers in rats after immediate sciatic nerve injury, with a decrease in muscle excitability (Gigo-Benato *et al.*, 2010).

Various methods have been used to apply PES both in vitro and in clinical practice (Fig. 1) (Balint et al., 2013; Griffin and Bayat, 2011). The simplest way to apply PES is to arrange that electrodes are in direct contact with cells cultured in vitro (Fig. 1a). However, some by-products such as ROS could be detrimental. Capacitive coupling for *in vitro* experiments consists in subjecting cells in a Petri dish to two parallel layers of electrode plates (Fig. 1b). Such an approach can generate a homogenous electrical field, when compared with direct coupling. To achieve a high field strength, a gap cuvette (Fig. 1c) was developed for in vitro experiments due to its short gap width of 2 or 4 mm. Such a cuvette can be used in electroporation for cell suspension and field strengths can be in kV/cm range. However, it cannot be used for repeated electrical stimulation because cells need to be suspended. In clinical practice, electrodes could be placed at the defect sites invasively, with a power source nearby (Fig. 1d), or could be placed outside the skin (Fig. 1e).

Cartilage regeneration and OA pose a formidable healthcare challenge. Interestingly, the fact that the application of PES, with appropriate parameters, could enhance regeneration of cartilage and ameliorate OA may open the door to potential clinical applications (Haddad et al., 2007). However, knowledge on the physical and cellular mechanisms of the biological effects of PES is still limited (Schoenbach et al., 2007). To make sense of the multitude and complexity of data within the scientific literature relating to PES and the observed beneficial therapeutic effects, the underlying cellular and molecular mechanisms associated with the application of PES were rigorously delineated, with a focus on chondrocytes, chondrogenic differentiation, and cartilage as models.

Physical models of the effects of electrical stimulation on cells

Conformational changes to biomolecules induced by electric fields elicit further biological responses. Several physical models for electrical stimulation of living cells, using varying cell parameters and electric



نے	
H	
·Ħ	
Ħ	
- 14-	
В	
5	
×	
e	
0	
č,	
· H	
2	
e	
σ	
C	
F	
· H	
Ħ	
4	
· 🔁	
5	
<u>p</u>	
В	
3	
7	
<u> </u>	
rti	
Ξ	
Ц	
T	
-	
<,	
щ	
еF	
he F	
the F	
y the F	
by the F	
l by the F	
d by the F	
red by the F	
ved by the F	
oved by the F	
roved by the F	
proved by the F	
Ipproved by the F	
approved by the F	
S approved by the F	
ES approved by the F	
PES approved by the F	
PES approved by the F	
rr PES approved by the F	
for PES approved by the F	
for PES approved by the F	
s for PES approved by the F	
ces for PES approved by the F	
ices for PES approved by the F	
vices for PES approved by the F	
evices for PES approved by the F	
Devices for PES approved by the F	
Devices for PES approved by the F	
1. Devices for PES approved by the F	
1. Devices for PES approved by the F	
le 1. Devices for PES approved by the F	
ble 1. Devices for PES approved by the F	
able 1. Devices for PES approved by the F	
Table 1. Devices for PES approved by the F	

				Parameters			Imnlanted	Product
Names of devices	Anatomical locations	Conditions	Frequency	Pulse duration	Intensity	Device class	device	code
Bladder system	Urinary bladder	Urination	7 or 9 MHz			HDE	yes	PAT
Cochlear implant with combined electrical stimulation and acoustic amplification	Ear	Hearing		20-400 µs		σ	Yes	PGQ
Dorsal root ganglion stimulator for pain relief	Dorsal root ganglion	Pain	4-80 Hz	40-1000 µs	4.6 V	σ	Yes	PMP
Intended to evaluate the functionality of human pain reception and transmission of sensory pathways	Wrist	Tremor symptom	$150\mathrm{Hz}$			7	No	NTU
Implanted phrenic nerve stimulator for central sleep apnea	Phrenic nerve	Central sleep apnea	20 Hz	60/150/300 µs		3	Yes	PSR
External vagal nerve stimulator for headache	Neck	Headache	5000 Hz		24 V, 60 mA	2	No	PKR
Limited output transcutaneous piezoelectric stimulator for skin reactions associated with insect bites	Skin	Pain		30 ms	222.83 mA	2	No	OSG
External pacemaker pulse generator	Right atrium and right ventricle	Heart disease				7	No	ĺΛΟ
Stimulator, electrical, implanted, for essential tremor	Deep brain structures	Tremor	$130\mathrm{Hz}$	60 ms		ß	Yes	МНΥ
Stimulator, electrical, implanted, for parkinsonian symptoms	Internal globus pallidus (gpi) or the subthalamic nucleus (STN)	Parkinson				ω	Yes	IHN
Transcutaneous electrical nerve stimulator for pain relief	Joint	Arthritis	2-100 Hz	50~250 µs	0.2~0.5 mA/ cm ²	7	No	OCF
Transcutaneous electrical nerve stimulator for pain relief	Skin	Pain		500 µs		2	No	NUH
Stimulator, low electric field, tumor treatment	Head	Tumor	100-300 kHz		0.7 V/cm	ю	No	NZK
Powered muscle stimulator	Skin	Muscle conditioning	1-120 Hz	340 µs		2	No	NGX
Transcutaneous electrical nerve stimulator to treat headache	Skin	Headache	1-100 Hz	0-250 µs		2	No	PCC
Transcutaneous electrical nerve stimulator for pain relief	Skin	Pain	$100\mathrm{Hz}$	100 µs		2	No	GZJ





Fig. 1. Device models for PES *in vitro* **and** *in vivo*. (**a**) Directly applied in a Petri dish *in vitro*. (**b**) Capacitive coupling in a Petri dish *in vitro*. (**c**) Gap cuvette. (**d**) Drect current *in vivo*. (**e**) Capacitive coupling *in vivo*.

fields, have been established. Physical models of membrane potential were established as early as 1953 when elliptical cells were exposed to direct electrical currents (Fricke, 1953), while the Schwan equation, based on models of spherical cells, represents one step forward by including extracellular fluid, plasma membrane, and cytoplasmic parameters (Schwan, 1956). The Schwan equation is as follows:

$\Delta \psi = 1.5 E_{\circ} R \cos \varphi$

where R is the cell radius, φ is the angle between the position vector and the applied electric field, E_0 is the field strength of the applied electric field, and $\Delta \psi$ is the change in membrane potential.

The multilayer dielectric model of spherical cells lays the foundation for a deeper understanding of the biological effects of PES, which include physical parameters of cell electrical conductivity, relative permittivity, cellular radius, and membrane thickness, as well as cell various organelles and their individual membrane thickness. Based on the multilayer dielectric model of the spherical cell, several conclusions can be drawn (Yao *et al.*, 2009):

1. membrane potentials are positively correlated with the field density of PES;

2. pores in the plasma membrane are formed within a transmembrane voltage range of about 0.5-1 V;

3. the effects of PES with a duration change of s to ps shift from plasma membranes to intracellular structures.

This, in turn, induces complex biological effects, such as activation or inhibition of signaling pathways, phosphorylation of proteins, manipulation of cell phenotypes, while longer durations or higher field strengths will induce apoptosis and necrosis (Weaver *et al.*, 2012) (Fig. 2). The upper boundary of physical parameters is mainly limited by thermal effects and cell death, while the lower boundary is limited by biological effects. Although duration and field density are two key parameters, based on these models and corresponding equations, frequency and timing of electrical stimulation are also critical due to their effect on the action area of PES (Weinberg, 2013). PES with a high frequency (10^4 - 10^9 Hz) would affect the inner membranes, while PES with a low frequency (< 10^4 Hz) may affect the plasma membrane (Yao *et al.*, 2009).

Field strength, duration, frequency, and time of stimulation are fundamental physical characteristics of PES. Field strength is the intensity of PES and it may range from mV/cm, V/cm, kV/cm to MV/cm for different types of application and studies. Duration of PES ranges from ps, ns, µs, m to s. Long duration plus high voltage would cause thermal effects. Frequency is an important parameter for PES, as PES with a high frequency (above 10 kHz) could affect intracellular organelles. The time span of stimulation is also important, which may be minutes to hours daily. Repeated stimulation would cause significantly different effects due to cumulative effects. For example, a pulse-number-dependent downregulation of mitochondrial function and cell numbers was observed by Hall et al. (2007) for nsPEFs. 3 d of PES would not cause significant cell death, while 7 d of PES would cause significant cell death (Kwon *et al.*, 2016). The formation of pores caused by PES increases due to the increase in pulse number (Yogesh, 2016). On the other hand, PES with a low field strength may require several hours daily to produce significant biological effects (Table 2).

The aforementioned physical model only works well when all individual cells have exactly the same physical properties, as well as disperse uniformly in the cell suspension (Yao *et al.*, 2009). However, distribution of the electric field on individual cells is not even, as cell suspension may not have optimal homogeneity while tissues are absolutely nonhomogenous. The effects of PES are also dependent on cell attachment, as well as changes in cell morphology (Casciola *et al.*, 2017). The distribution of PES is difficult to measure, but could be simulated with commercially available software, such as HFSS





Fig. 2. Biological effects of specific PES parameters. PES can cause different biological effects depending on its parameters. PES with high field strength and long duration can cause thermal effects and cell death. Reversible pore formation can be induced by PES with low field strength and short duration. Irreversible pore formation can be induced by PES with high field strength and long duration.

ANSYS, release 15.0 (ANSYS, Canonsburg, PA, USA) and COMSOL Multi-physics (COMSOL, Burlington, VT, USA) (Buyong *et al.*, 2015; Casciola *et al.*, 2017).

Effects of electrical stimulation on cells and their microenvironment

The effects of PES on different cellular components have been explored extensively, including VSDs, plasma membrane, intracellular components, and signaling pathways, which are often interwoven. PES induces conformational changes of voltagesensitive proteins, pore formation in the membrane, and calcium influx within seconds of its application, which, in turn, leads to activation of signaling pathways, cell migration, and cell death within minutes or hours.

VSDs

Conformational changes of proteins are early stage key biological effects of PES which are still unexplored. VSDs, key components of voltagegated channels, usually are comprised of four transmembrane helices (S1-S4) with a pore domain (helices S5 and S6). The S4 segment of voltage-gated channels with the motif [RK]-X(2)-R-X(2)-R-X(2)-[RK] is sensitive to electric fields (Murata et al., 2005). The basic residues of S4 segment are rapidly changed from hyperpolarized potentials, with an internally connected aqueous crevice, to depolarized potentials, with an externally connected aqueous crevice (Zhang et al., 2012). Electrical stimulation induces high levels of intracellular Na⁺ ions by the activation of Na/K ATPase, which is responsible for at least 35 % of the voltage-induced pores in the plasma membrane (Teissie and Tsong, 1980). However, electrical stimulation with a high intensity induces a negative-shift of channel open-threshold and reduces conductance of K⁺ channels, which may be due to conformational changes causing supraphysiological voltage in the membrane proteins of channel-gating systems (Chen, 2004). Furthermore, PES could denature voltage-sensitive membrane proteins with charged groups in some of their component amino acids which are sensitive to the plasma membrane potential (Chen, 2005). This may be due to breakage of chemical bonds between amino acids in the voltage-dependent membrane proteins, thus reducing their ability to open potassium ion channels simultaneously. From another point of view, nsPEFs inhibit voltage-gated Ca²⁺ and Na⁺ channels with reduction of the transmembrane ion gradient due to an increase in non-inactivating "leak" current (Nesin et al., 2012). Channel subtypes should also be considered, as voltage-gated potassium channels respond to electrical stimulation variably, depending on channel kinetics (Cameron et al., 2017). Reduction in the opening process of ion channels is less reversible in comparison to the process of inducing reversible pores by electroporation, which reclose within µs or ms. This mechanism is clinically significant as it helps illustrate how injuries caused by electric shock often show no signs of tissue damage but still result in malfunction (Chen, 2005). The presence of naturally occurring proteins with magnetic response hint at a possibility that there may be more proteins that have intrinsic capacity to respond to electrical stimulation (Qin et al., 2016).

Membrane

PES with varying durations has selective effects on either the plasma membrane or intracellular membranes. In terms of duration of pulses, PES is classified into several groups, including μ s, sub- μ s, ns, and ps (Yao *et al.*, 2009). Living cells respond to PES variably since they are conceptualized as a combination of capacitors and resistors with



Model	Electric parameter	Results	References
Fetal bovine articular chondrocytes	Frequency: 60 Hz Time: 0.5 h Intensity: 20 mV/cm	COLII and AGG \uparrow PG and collagen content \uparrow	(Wang <i>et al.,</i> 2004)
Adult bovine articular chondrocytes	Frequency: 60 kHz Intensity: 20 mV/cm Time: 4, 6 or 22 h	COLII and AGG ↑	(Xu et al., 2009)
Pig chondrocytes	Frequency: 1 Hz Pulse: 5; pulse duration: 100 ns Intensity: 10 kV/cm or 20 kV/cm	Dedifferentiation $COLI \uparrow, AGG \downarrow, COLII \downarrow, and$ $SOX9 \downarrow$	(Zhang <i>et al.,</i> 2014)
Human chondrocytes	Frequency: 4150 Hz Time: 30 min Intensity: 0.1 to 10 mV/cm Pulse duration: 10 ms	Proliferation ↑	(Fitzsimmons <i>et al.,</i> 2008)
Human chondrocytes	Frequency: 10 Hz Current: 10 μA Time: 3 or 6 h	Chondrocyte adhesion ↑ Long-term cell densities ↑	(Khang <i>et al.,</i> 2008)
Bovine articular chondrocytes	Frequency: 60 kHz Intensity: 0-45 mV/cm Time: 24 h	15-30 mV/cm: proliferation ↑ 45 mV/cm: proliferation ↓ and proteoglycan ↑	(Armstrong et al., 1988)

Table 2. Effects of PES on chondrocytes.

multilayer dielectric models (Schoenbach *et al.*, 2007; Yao *et al.*, 2009). Charging time constant is the time required to charge a plasma membrane (Deng *et al.*, 2003). The equations for charging time (τ) of the outer membrane and the inner membrane are as follows (Yao *et al.*, 2009). According to the multilayer dielectric model of a spherical cell, the constant τ_{cell} of the outer membrane (plasma membrane) can be expressed as

$$\tau_{cell} = \left(\frac{1}{2\gamma_o} + \frac{1}{\gamma_c}\right) \frac{\varepsilon_i \varepsilon_m}{d_m} R_c$$

The constant τ_{nuc} of the inner membrane (organelle membranes) can be expressed as

$$\tau_{nuc} = \left(\frac{1}{2\gamma_c} + \frac{1}{\gamma_{nc}}\right) \frac{\varepsilon_i \varepsilon_{nm}}{d_n} R_n$$

 $\gamma_{o'} \gamma_c$ and γ_{nc} are the conductivities of extracellular medium, cytoplasm and organelle cytoplasm, respectively. $\varepsilon_{i'} \varepsilon_m$ and ε_{nm} are the permittivities of vacuum, plasma membrane and organelle membrane, respectively. d_m and d_n denotes the thicknesses of the cell and organelle membranes, respectively. R_c and R_n are the radii of the cell and organelle, respectively. According to the above equations, the transmembrane potential of the membrane is as follows. At the end of the pulse, the outer membrane transmembrane potential is

$$V_m = 1.5 R_c E \left(1 - e^{-\frac{\tau}{\tau_{cell}}} \right) cos\theta$$

and the inner membrane transmembrane potential is

$$V_n = \frac{1.5\tau_{cell}R_nE}{\tau_{cell} - \tau_{nuc}} \left(e^{-\frac{\tau}{\tau_{cell}}} - e^{-\frac{\tau}{\tau_{nuc}}}\right)\cos\theta$$

E is the field intensity of the external electric field. θ is the polar angle measured with respect to the direction of the field. τ is the duration of the electric fields. Voltage of the plasma and organelle membranes are affected by the relationship of the size of τ , τ_{cell} and τ_{nuc} . τ_{cell} is in the range of hundreds of ns, while τ_{nuc} is in the range of about tens of ns. τ may be different for different cells, because different cells in different environment have different physiological parameters. If τ is much larger than $\tau_{cell'}$ then the outer membrane can be fully charged by electric fields. If τ is much smaller than $\tau_{cell'}$ then the outer membrane is poorly charged by electric fields. That is the same for the inner membrane. τ_{cell} is in the range of hundreds of ns for most cells and τ_{nuc} is in the range of tens of ns for most cells.

Based on the charge time of the plasma membrane, PES with durations of µs or longer, mainly exert effects on the cell outer membrane, while PES with a sub-us duration exert effects on both the inner and outer membrane with a sufficient PES intensity (at the kV/cm level) (Yao et al., 2009). nsPEFs with intensities up to 300 kV/cm have better penetrative capacity and mainly affect the inner cell membrane (Chopinet and Rols, 2015; Schoenbach, 2018). Theoretically, electrical stimulation with even shorter durations of ps could be endowed with even higher voltages. Ps-pulsed electrical stimulation with an intensity lower than 1 kV/cm hardly affects the plasma membrane (Yao et al., 2009). The intensity of electric fields is limited to less than 1 kV/cm, while the duration ranges from µs to ms (Zimmermann, 1986). Relatively short durations and high field strengths (hundreds kV/ cm) can potentially affect the intracellular membrane while restricting total energy and narrowing down the previously broad biological effects. In any case, precise control of PES is essential for various potential clinical applications and more attention should be paid to nsPEFs.

PES has the unique ability to induce reversible pore formation in plasma membranes within seconds,



including the outer plasma membrane, as well as intracellular organelles, such as mitochondria, endoplasmic reticulum, and nuclear membranes. PES induces changes in membranes immediately. Changes are reversible under PES with a low voltage and short duration, while irreversible under PES with a high voltage and long duration (Pakhomov *et al.*, 2015). Pores that form on plasma membranes are temporary and reversible when an electric field strength of about 0.1-20 kV/cm and pulse durations of 10 ns-10 ms are applied (Kirawanich et al., 2010; Kotnik et al., 2015). These temporary pores have been widely used in biomedical applications, including delivery of biomolecules, such as DNA, RNA, and proteins - commonly referred to as electroporation or electrofusion (Cemazar et al., 2013; Schoellhammer et al., 2014) - as well as drugs (Okino and Mohri, 1987). The voltage applied in electroporation can range between 200 V and 350 V, with 260 V being the most widely used (Nature Methods, 2006). The electrofusion technique is based on electroporation at 300 V pulse of the plasma membrane and is mediated by PEG (Hui and Stenger, 1993; Yu et al., 2008). The effects of electrofusion on different cell types, cell sizes, and intensity of electric field have been studied. Traditionally, the duration of electric fields used for electrofusion range between 10 and 100 µs (Jordan et al., 2013; Rems et al., 2013), while nsPEFs increase efficiency of cell electrofusion through electroporation of the contact areas between cells (Rems et al., 2014). Size of pores created by nsPEFs on the membrane are smaller than that caused by PES with longer durations (Vasilkoski et al., 2006).

Intracellular components and organelles

PES affects intracellular components and organelles, such as the cytoskeleton. PES increases speed of cell migration of both inner and outer meniscus cells (Yuan *et al.*, 2014) and ROS levels in both extracellular (electrochemical) and intracellular compartments (Pakhomova *et al.*, 2012). PES [10 ns, high voltages (> 150 kV/cm)] disrupts pre-messenger RNAs by inducing nuclear speckles of snRNPs (Chen *et al.*, 2007). PES with a low field strength (2.5 V/cm) accelerates cell migration within hours (Hayashi *et al.*, 2016). PES with a duration of ns can induce disassembly of actin structures and cell swelling resulting from cell permeabilization (Pakhomov *et al.*, 2014). Whereas, PES effects on ribosomes remain unexplored.

PES can induce apoptosis through dysfunctioning mitochondria. PES with a high field strength (kV/ cm) and a short duration of ns range decreases mitochondria membrane potential (Beebe *et al.*, 2012). The loss of mitochondria membrane potential may be a consequence of increased inner mitochondrial membrane permeability (Batista Napotnik *et al.*, 2012). PES with a duration of ps can induce the release of cytochrome C from mitochondria, with activation of caspase 3 and 9, resulting in apoptosis (Hua *et al.*, 2012).

PES also affects epigenetics through acetylation, histone modifications, alteration of the structural organization of chromatin, demethylation, as well as affecting RNA. PES downregulates histone deacetylase activities in HL-1 cells, with a significant reduction in Cx43 expression and cellcell communication (Meraviglia *et al.*, 2015). PES can also reversibly downregulate the activity of histone 1 kinase in pig oocytes (Leal and Liu, 1998). PES can prevent the tumorigenic transition of embryonic stem cells through downregulation of phospho-H3 and Ki67 immuno-reactivity (Yamada *et al.*, 2007). PES promotes global DNA demethylation and downregulates *DNMT1* expression (unpublished data).

Signaling pathways

PES exerts effects on cells through activation or inhibition of various signaling pathways (Liu and Song, 2014). PES activates protein kinase C signaling pathway by inducing PIP2 hydrolysis (Tolstykh et al., 2013). PES inhibits VEGF signaling pathway via downregulation of VEGF and PD-ECGF, thereby disrupting pro-angiogenic and anti-angiogenic balance, which may consequently inhibit cancer development and suppress tumor blood vessel growth (Chen et al., 2007). PES activates MAPK signaling pathway. PES not only upregulates phosphorylation of ERK and p38 (Zhao et al., 2006), but can also activate JNK, another member of the MAPK family, with fast increased phosphorylation level within minutes, which then decreases within 1 h (Morotomi-Yano et al., 2011a; 2011b). As the signaling pathways activated by PES may become inactivated within minutes or hours, the effect caused by PES may last for minutes, hours, or days. PESpreconditioning promotes cartilage regeneration within weeks through enhancing differentiation of MSCs (Ning et al., 2019). This is the consequence of the biological effects initiated by PES during the early stage after stimulation. Meanwhile, phosphorylation of MAPK caused by PES is reversible (Morotomi-Yano et al., 2011a). PES can also induce dedifferentiation of chondrocytes through activation of Wnt/β-catenin signaling pathway (Zhang et al., 2014). Activation of β -catenin signaling pathway enhances NEUROG2 expression in brain injury after PES treatment (Matsumoto *et al.*, 2013) (Fig. 3).

PES exerts effects on calcium-related signaling pathways through the release of Ca²⁺ ions from endoplasmic reticulum compartments (Semenov *et al.*, 2013; Vernier *et al.*, 2003). PES with high field strength can induce calcium release within seconds (Zhang *et al.*, 2014). PES has been reported to induce release of Ca²⁺ leading to an increase in cytoskeletal CaM levels (Brighton *et al.*, 2001). PES can also induce calcium influx from the external medium (Bourguignon *et al.*, 1989). PES enhances proliferation of cardiac fibroblast and myocardial fibrosis through activation of the CaN-NFAT pathway, regulated by Ca²⁺/CaM (Chen *et al.*, 2012). PES increases synthesis



of NO and production of cGMP through calcium/ CaM and inhibits Ca²⁺/CaM, thereby blocking these effects (Fitzsimmons *et al.*, 2008).

PES can also regulate the process of apoptosis and its associated biochemical markers. PES activates caspase-dependent signaling pathways in apoptosis, with the activation of caspase-12, -9, and -3, leading to release of cytochrome c, upregulation of Bax and downregulation of Bcl-2 in HeLa cells (Chen *et al.*, 2013).

Physiological microenvironment

PES can alter blood flow, temperature, and pH of tissues. Blood flow is significantly increased following application of electrical stimulation in wound healing (Ud-Din et al., 2015). Increased blood flow following increasing capillary density is observed when the transcutaneous oxygen partial pressure is increased by 82.4 % during treatment of CVI (Jünger et al., 1997). Thermal effects are integral in traditional PES, but on the other hand, thermal effects are dependent on tissue properties as well as stimulation parameters (*i.e.*, frequency, pulse duration, field strength) (Schoenbach et al., 2004). Electrical stimulation increases temperature around the electrodes in the visual prosthesis system (Çelik and Karagöz, 2014). Temperature is increased around the electrodes (both anode and cathode) and the pH value varied from 3 (near the anode) to 12 (near the cathode) in the electrode array for local control of solid tumors (Soba et al., 2018). Low-voltage electrical stimulation can promote muscle tenderness and accelerate glycolysis, with a significant fall in pH during the first 6 h post mortem (Polidori et al., 1999).

PES has been reported to cause deleterious effects on cells, including cell death and cell fragmentation (Yang *et al.*, 2018). PES with high field strength and long duration can result in thermal effects due to Joule heating (Yarmush *et al.*, 2014), with consequent cell death, as proteins denature when the temperature reaches about 43-45 °C. Also, nsPESs with high field strength lead to cell death within minutes, which is mainly due to cell fragmentation (Ning *et al.*, 2019). PES with high field strength and long duration can also induce cell death, following both necrotic and apoptotic ways (Pakhomova *et al.*, 2013).

Effects of PES on chondrocytes, mesenchymal stem cells, cartilage explants, and OA

PES promotes cartilage regeneration and ameliorate OA through modulation of extracellular matrix and various biological factors, such as BMPs, TGF- β , and IGF2, which in turn induce anti-inflammatory and anabolic effects to improve articular cartilage regeneration together with amelioration of inflammation (Haddad et al., 2007; Massari et al., 2007). Cell-based repair strategies induce cartilage regeneration (Johnstone et al., 2013). Chondrocytes and MSCs (or their chondrogenic progenies) are important cell sources (de Vries-van Melle et al., 2014; Estes et al., 2010; Guilak et al., 2010). Both ADSCs and BMSCs have demonstrated their unique responses to growth-factor-induced chondrogenic differentiation, respectively to TGF-B3 and BMP6 (Diekman et al., 2009). However, there are obstacles to using primary chondrocytes for cell culture applications, including limited cell numbers and dedifferentiated phenotypes during in vitro expansion (Chung and Burdick, 2008). Challenges faced in culturing bone-marrow-derived MSCs or ADSCs in vitro include low efficiency of chondrogenic differentiation and heterogeneous phenotypes (Perdisa et al., 2015; Somoza et al., 2014), while proper application of PES could potentially overcome these obstacles.

Chondrocytes

PES can induce varied or even opposite effects on chondrocytes, including increased proliferation and increased or decreased differentiation (represented as expression of *ColII* or ratio of *ColII/ColI* and GAG production). PES for 24 h at 15-30 mV/cm increases proliferation of primary chondrocytes harvested



Fig. 3. Signalling pathways activated by PES for a generic cell. MAPK and its downstream signalling pathways are activated by PES. Calcium influx is induced by PES. Cell death could be induced by PES through the caspase signalling pathway. Wnt/ β -catenin signalling pathway could be activated by PES.



Model	Other factors	Electrical parameters	Results	References
Calf cartilage disks		Frequency: 1, 10, 100, 10 ³ , and 10 ⁴ Hz Intensity: 10-30 mA/cm ²	Protein synthesis ↑	(Macginitie <i>et al.,</i> 1994)
Bovine articular cartilage explants		Frequency: 60 Hz Intensity: 20 mV/cm Time: 1 h on and 5 h off, 4 times/d for 3 d	Production of proteoglycan and collagen ↑ AGG and COLII ↑	(Brighton <i>et al.,</i> 2006)
Human osteoarthritic cartilage explants	With or without IL-1β	Frequency: 60 kHz Intensity: 20 mV/cm Time: 1 h on and 5 h off, 4 times/d for 3 d	With IL-1β: production of proteoglycan and collagen ↑ AGG and COLII ↑	(Brighton <i>et al.,</i> 2008)

Table 3. Effects of PES on cartilage explants.

from growth plate cartilage, with upregulated [³H] thymidine uptake, and enhances proteoglycan deposition, with upregulated [35S] sulfate uptake, while PES for 24 h at 45 mV/cm inhibits proliferation and enhances proteoglycan deposition (Armstrong et al., 1988). PES (0.5 h, 20 mV/cm, 60 Hz) can increase COLII and AGG expression in fetal bovine chondrocytes in a micromass culture system (Wang et al., 2004). PES (4, 6 or 22 h, 60 kHz, 20 mV/cm) induces up-regulation of both COLII and AGG expression levels through extracellular Ca²⁺ influx through voltage-gated calcium channels rather than intracellular Ca²⁺ repositories (Xu *et al.*, 2009). PES (3 or 6 h, 10 Hz, 10 µA) enhances chondrocyte adhesion and long-term cell densities for up to 2 d (Khang et al., 2008). On the other hand, nsPEFs (5 pulses, 1 Hz, 100 ns at 10 or 20 kV/cm) decreases gene expression levels of COLII, AGG and SOX9 through activation of Wnt/\beta-catenin signaling pathway and intracellular calcium ion efflux (Zhang et al., 2014) (Table 2).

This obvious inconsistency could be attributed to either varying parameters of PES or cell types with ever-changing intra- and inter-cellular structures. Generally, PES with low electric field strength and long duration promotes proliferation with effects mainly on calcium homeostasis, cell skeleton, or membrane, while PES with high electric field strength affects intracellular structures or more specific targets.

Chondrogenic differentiation of MSCs

PES has been optimized to promote chondrogenic differentiation of MSCs (Table 4). Piezoelectric scaffolds, exhibiting low voltage output, enhance chondrogenic differentiation of MSCs with upregulation of intense proteoglycan staining and *COLII*, while piezoelectric scaffolds with high voltage output enhance osteogenic differentiation (Damaraju *et al.*, 2017). This emphasizes the importance of PES parameters in chondrogenic differentiation. PES (20 min daily for 7 d, 20 mV/cm, 60 KHz), with or without TGF- β 3, promotes chondrogenic differentiation, with enhanced *COLII* and *SOX9* expression levels (Esfandiari *et al.*, 2014). PES (20 min, 1 kHz, 20 mV/cm) leads to chondrogenic

differentiation of ADSCs (Mardani et al., 2016). TGF- β , BMPs and other signaling pathways have been implicated in PES treatment. PES (1, 5 or 25 V/ cm, 5 Hz) promotes chondrogenic differentiation via upregulated SOX9, COLII, and AGG, with TGF- β signaling, BMP signaling and extracellular ATP signaling via P2X₄ receptors (Kwon et al., 2016). Electrical stimulation induces calcium oscillations through modulation of VOCCs, which are implicated in chondrogenesis (Uzieliene et al., 2018). PES (nsscale with high intensity) enhances chondrogenic differentiation of MSCs with significantly upregulated expression levels of SOX9, COLII, and AGG, together with down-regulated expression of COLX through activation of c-Jun and Stat3 signaling pathways (Ning et al., 2019). ns-PESs promote chondrogenic differentiation of MSCs through activation of MAPK signaling pathways (Morotomi-Yano et al., 2011a).

PES can promote chondrogenic differentiation through upregulation of related growth factors. PES (50 and 100 mV/mm) also increases FGF2 secretion and upregulates phosphorylation levels of Smad2 and Smad3 in fibroblasts (Wang et al., 2017). PES (50 mV/ mm) upregulates the secretion of FGF1 and FGF2 in fibroblasts (Rouabhia et al., 2013). PES (5 V, 0.1 ms, 10 min) together with a heat shock can activate insulin signaling through phosphorylation of IRS (Morino et al., 2008). However, disruption of the secondary structure of insulin is observed at higher electric field strengths (above 0.25 V/nm) within 1 µs (Wang et al., 2014) and the binding capacity of insulin to receptors is significantly reduced by long PES duration (0.7 V/m, 50 Hz, 20 min) (Li et al., 2005). So, whether insulin is involved in chondrogenic differentiation of MSCs by PES remains to be explored. Perhaps low intensity PES can activate insulin signaling pathway to promote chondrogenic differentiation while high intensity PES can not.

Cartilage explants and OA

PES can increase matrix production and reduce matrix destruction in cartilage explants. PES (60 kHz, 20 mV/cm) increases production of proteoglycan and collagen in the absence or presence of IL-1 β , while dramatically inhibiting expression of matrix



Model	Electrical parameter	Results	References	
Piezoelectric scaffolds Human MSCs	20 mV/mm	GAG and COLII ↑	(Damaraju <i>et al.,</i> 2017)	
Human ADSCs	Frequency: 60 KHz Intensity: 20 mV/cm Time: 20 min daily for 7 d	COLII and SOX9↑	(Esfandiari <i>et al.,</i> 2014)	
ADSCs	Frequency: 1 kHz Intensity: 20 mv/cm Time: 20 min COLII and SOX9↑		(Mardani <i>et al.,</i> 2016)	
Mouse MSCs	Frequency: 5 Hz Intensity: 1, 5 or 25 V/cm	COLII and SOX9 ↑ COLI ↓	(Kwon <i>et al.,</i> 2016)	
Pig MSCs	Frequency: 1 Hz Intensity: 10 kV/cm, 20 kV/cm Duration: 10 ns, 100 ns	COLII and SOX9↑	(Ning et al., 2019)	

Table 4	. Effects	of PES	on the	chondrogenic	differentiation	of MSCs.
---------	-----------	--------	--------	--------------	-----------------	----------

metalloproteinase in full-thickness osteoarthritic adult human articular cartilage explants (Brighton *et al.*, 2008). Brighton *et al.* (2006) observed that PES (60 kHz, 20 mV/cm) increases collagen production and expression levels of *COLII* and *AGG* within adult bovine articular cartilage explants (Table 3).

PES used for neuromuscular stimulation increased muscle thickness and pennation angle in 45 women (age 66-75 years) with knee OA and it resulted in a significant increase in functional capacity, as measured by the 6 min walk test and Timed Up and Go Test (Melo Mde *et al.*, 2015). PES resulted in high global effectiveness of OA treatment in two patients (Fary *et al.*, 2009). However, the third patient exhibited no change, emphasizing the importance of heterogeneity. PES resulted in a significant downregulation of various OA symptoms in hands, with respect to pain, swelling, grip strength, and pinch force (Holt *et al.*, 2018).

The effects of PES on OA pain relief remains to be explored. Neuromuscular electrical stimulation reduces pain (de Oliveira Melo et al., 2016). PES can alleviate pain in OA. PES (at least 6 h/d for 3 months) significantly reduced pain in patients with knee OA in clinical trials (Garland et al., 2007). Although PES could improve physical functions in subjects with knee OA, it may not reduce pain. PES attenuated knee OA symptoms in a dose-response manner after 750 h in a clinical trial with 288 patients who had failed nonsurgical therapy (Farr *et al.*, 2006). However, contrary results were also reported in which PES could not provide symptomatic relief for OA (Fary et al., 2008; 2011). This difference may be due to limitations of the sample size and heterogeneity, such as age or BMI. Maybe the type of electrical stimulation could change the effects of PES on pain relief. Among electrical stimulation therapy, from h-TENS, l-TENS, NMES, IFC, PES to NIN, IFC seems to be the best treatment modality in terms of effectiveness, as compared with PES and other electrical stimulation modalities (Zeng *et al.*, 2015). Amplitude-modulated frequency generated by IFC can permeate more deeply, which could improve pain relief, as the main analgesic component of IFC (Johnson et al., 2003). Meanwhile,

mechanisms in the observed effects of electrical stimulation on OA need to be further explored to optimize the treatment, such as thermal effects, cell migration, blood flow *etc.*. Perhaps PES with a low frequency (≤ 100 Hz) is able to improve physical function but not pain intensity (Negm *et al.*, 2013), which calls for mechanistic exploration and studies using PES with high frequency and short duration or other parameters to avoid harmful changes.

Perspectives

Most research studies on PES were stalled after identification of specific signaling pathways involved, without elucidating the initiator molecule or initial conformational changes of biomolecules. Some studies have moved forward to identifying a complex array of signaling pathways involved, with the biological effects observed being just the outcome of multiple well-orchestrated processes. A comprehensive overview of biological mechanisms, as well as the corresponding physical stimuli involved, are needed to enable more extensive clinical applications by using novel toolkits. The target may be a protein subunit or some other molecular subunit that requires identification by high-throughput assays.

Integration with novel technology platforms is necessary to facilitate clinical applications of PES, *i.e.* high-throughput screening, microfluidics platforms, single-cell analysis, and bioinformatics. Novel technologies such as nsPEFs, which could precisely trigger specific activities within the cell and organelle internally, should be further investigated. PES with different parameters induces variable effects on different organelles, which in turn elicit variable biological effects on cells, tissues, and organs. The specific parameters of PES will depend on the clinical conditions of individual patients, different tissue states, and cell-types. Therefore, utilizing PES with appropriate parameters, such as specific stimulation frequencies and specific durations, must be optimized and standardized by high-throughput assays and www.ecmjournal.org





comprehensive biological mechanistic studies, to enable PES technology to be applied clinically for enhancing cartilage repair and regeneration. The diameter of chondrocytes is about 20 μ m, while that of MSCs is 10-15 μ m. This means that higher field strengths are required for chondrocyte than MSCs. Lower field strength (mv/cm), moderate duration (μ sms), and hours-long stimulation would be preferred for cartilage repair and chondrogenic differentiation of MSCs, as ns-pulsed electric fields with high field strength may cause dedifferentiation of chondrocytes (Table 2) (Zhang *et al.*, 2014).

Wnt/β-catenin, TGF-β, MAPK, and FGF signaling pathways are critical for cartilage repair and chondrogenic differentiation of MSCs (Tang et al., 2015) (Fig. 3). TGF- β /Smad signaling pathway is activated by PES during chondrogenesis of MSCs (Kwon et al., 2016). Wnt/β-catenin is activated in primary chondrocytes by nsPESs (Zhang et al., 2014). MAPK signaling pathway is activated in enhanced chondrogenic differentiation of MSCs induced by nsPEFs (Ning et al., 2019). PES (50 mV/mm) promotes secretion of FGF-1 and FGF-2 in fibroblasts (Rouabhia et al., 2013). TGF- β and FGF signaling pathways need to be further explored during chondrogenic differentiation of MSCs induced by PES, particularly for upstream and downstream signaling pathways. Possible signaling pathways need to be explored (Fig. 4).

Acknowledgement

Authors would like to thank Rossana Rico for revision of the manuscript and Prof Jue Zhang and Jinsong Guo for constructive suggestions. This work was supported by the China National Natural Science Foundation (81772334).

No competing financial interests exist.

There are no ethical issues pertaining to human and animal rights.

References

Armstrong PF, Brighton CT, Star AM (1988) Capacitively coupled electrical stimulation of bovine growth plate chondrocytes grown in pellet form. J Orthop Res **6**: 265-271.

Ashley Z, Salmons S, Boncompagni S, Protasi F, Russold M, Lanmuller H, Mayr W, Sutherland H, Jarvis JC (2007) Effects of chronic electrical stimulation on long-term denervated muscles of the rabbit hind limb. J Muscle Res Cell Motil **28**: 203-217.

Balint R, Cassidy NJ, Cartmell SH (2013) Electrical stimulation: a novel tool for tissue engineering. Tissue Eng Part B Rev **19**: 48-57.

Batista Napotnik T, Wu YH, Gundersen MA, Miklavcic D, Vernier PT (2012) Nanosecond electric pulses cause mitochondrial membrane permeabilization in Jurkat cells. Bioelectromagnetics **33**: 257-264.

Beebe SJ, Chen Y-J, Sain NM, Schoenbach KH, Xiao S (2012) Transient features in nanosecond pulsed electric fields differentially modulate mitochondria and viability. PLoS One 7: e51349. DOI: 10.1371/ journal.pone.0051349.

Bourguignon GJ, Jy W, Bourguignon LYW (1989) Electric stimulation of human fibroblasts causes an increase in Ca2+ influx and the exposure of additional insulin receptors. J Cell Physiol **140**: 379-385.

Brighton CT, Wang W, Clark CC (2008) The effect of electrical fields on gene and protein expression in



human osteoarthritic cartilage explants. J Bone Joint Surg Am **90**: 833-848.

Brighton CT, Wang W, Clark CC (2006) Upregulation of matrix in bovine articular cartilage explants by electric fields. Biochem Biophys Res Commun **342**: 556-561.

Brighton CT, Wang W, Seldes R, Zhang GH, Pollack SR (2001) Signal transduction in electrically stimulated bone cells. J Bone Joint Surg Am **83**: 1514-1523.

Buyong MR, Yunas J, Hamzah AA, Majlis BY, Larki F, Aziz NA (2015) Design, fabrication and characterization of dielectrophoretic microelectrode array for particle capture. Microelectronics International **32**: 96-102.

Cameron MA, Al Abed A, Buskila Y, Dokos S, Lovell NH, Morley JW (2017) Differential effect of brief electrical stimulation on voltage-gated potassium channels. J Neurophysiol **117**: 2014-2024.

Casciola M, Xiao S, Pakhomov AG (2017) Damagefree peripheral nerve stimulation by 12-ns pulsed electric field. Sci Rep 7: 10453. DOI: 10.1038/s41598-017-10282-5.

Çelik ME, Karagöz İ (2014) The effect of the electrical stimulation on temperature rise in the retinal tissue for visual prostheses. IJCEE **6**: 369-372.

Cemazar J, Miklavcic D, Kotnik T (2013) Microfluidic devices for manipulation, modification and characterization of biological cells in electric fields – a review. Informacije Midem-Ljubljana **43**: 143-161.

Chao E, Inoue N (2003) Biophysical stimulation of bone fracture repair, regeneration and remodelling. Eur Cell Mater **6**: 72-84.

Chen QQ, Zhang W, Chen XF, Bao YJ, Wang J, Zhu WZ (2012) Electrical field stimulation induces cardiac fibroblast proliferation through the calcineurin-NFAT pathway. Can J Physiol Pharmacol **90**: 1611-1622.

Chen WJ, Xiong ZA, Zhang M, Yao CG, Zhao ZY, Hua YY, Zhou W (2013) Picosecond pulsed electric fields induce apoptosis in HeLa cells *via* the endoplasmic reticulum stress and caspase-dependent signaling pathways. Int J Oncol **42**: 963-970.

Chen W (2005) Electroconformational denaturation of membrane proteins. Ann N Y Acad Sci **1066**: 92-105.

Chen W (2004) Supra-physiological membrane potential induced conformational changes in K+ channel conducting system of skeletal muscle fibers. Bioelectrochemistry **62**: 47-56.

Chopinet L, Rols M-P (2015) Nanosecond electric pulses: a mini-review of the present state of the art. Bioelectrochemistry **103**: 2-6.

Chung C, Burdick JA (2008) Engineering cartilage tissue. Adv Drug Deliv Rev **60**: 243-262.

Ciombor DM, Aaron RK (2005) The role of electrical stimulation in bone repair. Foot Ankle Clin **10**: 579-593.

Damaraju SM, Shen Y, Elele E, Khusid B, Eshghinejad A, Li J, Jaffe M, Arinzeh TL (2017) Threedimensional piezoelectric fibrous scaffolds selectively promote mesenchymal stem cell differentiation. Biomaterials **149**: 51-62.

de Oliveira Melo M, Pompeo KD, Baroni BM, Vaz MA (2016) Effects of neuromuscular electrical stimulation and low-level laser therapy on neuromuscular parameters and health status in elderly women with knee osteoarthritis: a randomized trial. J Rehabil Med **48**: 293-299.

de Vries-van Melle ML, Tihaya MS, Kops N, Koevoet W, Murphy JM, Verhaar J, Alini M, Eglin D, van Osch G (2014) Chondrogenic differentiation of human bone marrow-derived mesenchymal stem cells in a simulated osteochondral environment is hydrogel dependent. Eur Cell Mater **27**: 112-123.

Deng J, Schoenbach KH, Buescher ES, Hair PS, Fox PM, Beebe SJ (2003) The effects of intense submicrosecond electrical pulses on cells. Biophysical J 84: 2709-2714.

Diekman BO, Rowland CR, Lennon DP, Caplan AI, Guilak F (2009) Chondrogenesis of adult stem cells from adipose tissue and bone marrow: induction by growth factors and cartilage-derived matrix. Tissue Eng Part A **16**: 523-533.

Ebrahim S, Mollon B, Bance S, Busse JW, Bhandari M (2014) Low-intensity pulsed ultrasonography *versus* electrical stimulation for fracture healing: a systematic review and network meta-analysis. Can J Surg **57**: E105-118.

Esfandiari E, Roshankhah S, Mardani M, Hashemibeni B, Naghsh E, Kazemi M, Salahshoor M (2014) The effect of high frequency electric field on enhancement of chondrogenesis in human adiposederived stem cells. Iran J Basic Med Sci **17**: 571-576.

Estes BT, Diekman BO, Gimble JM, Guilak F (2010) Isolation of adipose-derived stem cells and their induction to a chondrogenic phenotype. Nat Protoc 5: 1294-1311.

Farr J, Mont MA, Garland D, Caldwell JR, Zizic TM (2006) Pulsed electrical stimulation in patients with osteoarthritis of the knee: follow up in 288 patients who had failed non-operative therapy. Surg Technol Int **15**: 227-233.

Fary RE, Briffa NK, Briffa TG (2009) Effectiveness of pulsed electrical stimulation in the management of osteoarthritis of the knee: three case reports. Physiother Theory Pract **25**: 21-29.

Fary RE, Carroll GJ, Briffa TG, Briffa NK (2011) The effectiveness of pulsed electrical stimulation in the management of osteoarthritis of the knee results of a double-blind, randomized, placebo-controlled, repeated-measures trial. Arthritis Rheum **63**: 1333-1342.

Fary RE, Carroll GJ, Briffa TG, Gupta R, Briffa NK (2008) The effectiveness of pulsed electrical stimulation (E-PES) in the management of osteoarthritis of the knee: a protocol for a randomised controlled trial. BMC Musculoskelet Disord **9**: 18. DOI: 10.1186/1471-2474-9-18.

Fitzsimmons RJ, Gordon SL, Kronberg J, Ganey T, Pilla AA (2008) A pulsing electric field (PEF)



increases human chondrocyte proliferation through a transduction pathway involving nitric oxide signaling. J Orthop Res **26**: 854-859.

Fricke H (1953) The electric permittivity of a dilute suspension of membrane-covered ellipsoids. J Appl Phys **24**: 644-646.

Garland D, Holt P, Harrington JT, Caldwell J, Zizic T, Cholewczynski J (2007) A 3-month, randomized, double-blind, placebo-controlled study to evaluate the safety and efficacy of a highly optimized, capacitively coupled, pulsed electrical stimulator in patients with osteoarthritis of the knee. Osteoarthritis Cartilage **15**: 630-637.

Gigo-Benato D, Russo TL, Geuna S, Santa Rosa Domingues NR, Salvini TF, Parizotto NA (2010) Electrical stimulation impairs early functional recovery and accentuates skeletal muscle atrophy after sciatic nerve crush injury in rats. Muscle Nerve **41**: 685-693.

Griffin M, Bayat A (2011) Electrical stimulation in bone healing: critical analysis by evaluating levels of evidence. Eplasty **11**: e34.

Guilak F, Estes BT, Diekman BO, Moutos FT, Gimble JM (2010) 2010 Nicolas Andry Award: multipotent adult stem cells from adipose tissue for musculoskeletal tissue engineering. Clin Orthop Relat Res **468**: 2530-2540.

Hall Eh, Schoenbach Kh, Beebe S (2007) Nanosecond pulsed electric fields induce apoptosis in p53-wildtype and p53-null HCT116 colon carcinoma cells. Apoptosis **12**:1721-1731.

Haddad JB, Obolensky AG, Shinnick P (2007) The biologic effects and the therapeutic mechanism of action of electric and electromagnetic field stimulation on bone and cartilage: new findings and a review of earlier work. J Altern Complement Med **13**: 485-490.

Hayashi H, Edin F, Li H, Liu W, Rask-Andersen H (2016) The effect of pulsed electric fields on the electrotactic migration of human neural progenitor cells through the involvement of intracellular calcium signaling. Brain Research **1652**: 195-203.

Hirt MN, Boeddinghaus J, Mitchell A, Schaaf S, Börnchen C, Müller C, Schulz H, Hubner N, Stenzig J, Stoehr A (2014) Functional improvement and maturation of rat and human engineered heart tissue by chronic electrical stimulation. J Mol Cell Cardiol **74**: 151-161.

Holt PA, Ozyurekoglu T, Deveshwar S, Maclaughlin EJ, Khlopas A, Mont MA, Pang SW, Tuber JS, Schechtman J, Zizic TM (2018) A pulsed electrical joint stimulator for the treatment of osteoarthritis of the hand and wrist. Orthopedics **41**: e550-e556.

Hua YY, Wang XS, Zhang Y, Yao CG, Zhang XM, Xiong ZA (2012) Intense picosecond pulsed electric fields induce apoptosis through a mitochondrialmediated pathway in HeLa cells. Mol Med Rep 5: 981-987.

Hui SW, Stenger DA (1993) Electrofusion of cells: hybridoma production by electrofusion and

polyethylene glycol. Methods in Enzymology **220**: 212-227.

Johnson MI, Tabasam G (2003) A single-blind investigation into the hypoalgesic effects of different swing patterns of interferential currents on coldinduced pain in healthy volunteers. Arch Phys Med Rehabil **84**: 350-357.

Johnstone B, Alini M, Cucchiarini M, Dodge GR, Eglin D, Guilak F, Madry H, Mata A, Mauck RL, Semino CE (2013) Tissue engineering for articular cartilage repair—the state of the art. Eur Cell Mater **25**: 248-256.

Jordan CA, Neumann E, Sowers AE (2013) Electroporation and electrofusion in cell biology. Springer Science & Business Media. New York, pp: 208-220.

Jünger M, Zuder D, Steins A, Hahn M, Klyscz T (1997) Treatment of venous ulcers with low frequency pulsed current (Dermapulse): effects on cutaneous microcirculation. Hautarzt **48**: 897-903.

Kern H, Barberi L, Lofler S (2016) Electrical stimulation counteracts muscle decline in seniors. Front Aging Neurosci 6: 189. DOI: 10.3389/fnagi.2014.00189.

Khang D, Park GE, Webster TJ (2008) Enhanced chondrocyte densities on carbon nanotube composites: the combined role of nanosurface roughness and electrical stimulation. J Biomed Mater Res A **86**: 253-260.

Kirawanich P, Pausawasdi N, Srisawat C, Yakura SJ, Islam NE (2010) An FDTD interaction scheme of a high-intensity nanosecond-pulsed electric-field system for *in vitro* cell apoptosis applications. IEEE **38**: 2574-2582.

Kotnik T, Frey W, Sack M, Meglič SH, Peterka M, Miklavčič D (2015) Electroporation-based applications in biotechnology. Trends Biotechnol **33**: 480-488.

Kwon HJ, Lee GS, Chun H (2016) Electrical stimulation drives chondrogenesis of mesenchymal stem cells in the absence of exogenous growth factors. Sci Rep **6**: 39302. DOI: 10.1038/srep39302.

Leal CL, Liu L (1998) Differential effects of kinase inhibitor and electrical stimulus on activation and histone H1 kinase activity in pig oocytes. Anim Reprod Sci **52**: 51-61.

Li L, Dai Y, Xia R, Chen S, Qiao D (2005) Pulsed electric field exposure of insulin induces anti-proliferative effects on human hepatocytes. Bioelectromagnetics **26**: 639-647.

Liu Q, Song B (2014) Electric field regulated signaling pathways. Int J Biochem Cell Biol **55**: 264-268.

Mardani M, Roshankhah S, Hashemibeni B, Salahshoor M, Naghsh E, Esfandiari E (2016) Induction of chondrogenic differentiation of human adipose-derived stem cells by low frequency electric field. Adv Biomed Res **5**: 97. DOI: 10.4103/2277-9175.183146.

Massari L, Benazzo F, De MM, Setti S, Fini M (2007) Effects of electrical physical stimuli on articular cartilage. J Bone Joint Surg Am **89** Suppl 3: 152-161.



Matsumoto M, Imura T, Fukazawa T, Sun Y, Takeda M, Kajiume T, Kawahara Y, Yuge L (2013) Electrical stimulation enhances neurogenin2 expression through beta-catenin signaling pathway of mouse bone marrow stromal cells and intensifies the effect of cell transplantation on brain injury. Neurosci Lett **533**: 71-76.

Melo Mde O, Pompeo KD, Brodt GA, Baroni BM, da Silva Junior DP, Vaz MA (2015) Effects of neuromuscular electrical stimulation and lowlevel laser therapy on the muscle architecture and functional capacity in elderly patients with knee osteoarthritis: a randomized controlled trial. Clin Rehabil **29**: 570-580.

Meraviglia V, Azzimato V, Colussi C, Florio MC, Binda A, Panariti A, Qanud K, Suffredini S, Gennaccaro L, Miragoli M (2015) Acetylation mediates Cx43 reduction caused by electrical stimulation. J Mol Cell Cardiol **87**: 54-64.

Morino S, Kondo T, Sasaki K, Adachi H, Suico MA, Sekimoto E, Matsuda T, Shuto T, Araki E, Kai H (2008) Mild electrical stimulation with heat shock ameliorates insulin resistance *via* enhanced insulin signaling. PloS One **3**: e4068. DOI: 10.1371/journal. pone.0004068.

Moro E, Esselink R, Xie J, Hommel M, Benabid A, Pollak P (2002) The impact on Parkinson's disease of electrical parameter settings in STN stimulation. Neurology **59**: 706-713.

Morotomi-Yano K, Akiyama H, Yano K (2011a) Nanosecond pulsed electric fields activate MAPK pathways in human cells. Arch Biochem Biophys **515**: 99-106.

Morotomi-Yano K, Uemura Y, Katsuki S, Akiyama H, Yano K (2011b) Activation of the JNK pathway by nanosecond pulsed electric fields. Biochem Biophys Res Commun **408**: 471-476.

Murata Y, Iwasaki H, Sasaki M, Inaba K, Okamura Y (2005) Phosphoinositide phosphatase activity coupled to an intrinsic voltage sensor. Nature **435**: 1239-1243.

Nature Methods (2006) Transfection of mammalian cells by electroporation. Nature Methods **3**: 67-68.

Negm A, Lorbergs A, Macintyre N (2013) Efficacy of low frequency pulsed subsensory threshold electrical stimulation *vs* placebo on pain and physical function in people with knee osteoarthritis: systematic review with meta-analysis. Osteoarthritis Cartilage **21**: 1281-1289.

Nesin V, Bowman AM, Xiao S, Pakhomov AG (2012) Cell permeabilization and inhibition of voltagegated Ca2+and Na+ channel currents by nanosecond pulsed electric field. Bioelectromagnetics **33**: 394-404.

Ning T, Guo J, Zhang K, Li K, Zhang J, Yang Z, Ge Z (2019) Nanosecond pulsed electric fields enhanced chondrogenic potential of mesenchymal stem cells *via* JNK/CREB-STAT3 signaling pathway. Stem Cell Res Ther **10**: 45. DOI: 10.1186/s13287-019-1133-0.

Okino M, Mohri H (1987) Effects of a high-voltage electrical impulse and an anticancer drug on *in vivo* growing tumors. Jpn J Cancer Res **78**: 1319-1321.

Pakhomov AG, Gianulis E, Vernier PT, Semenov I, Xiao S, Pakhomova ON (2015) Multiple nanosecond electric pulses increase the number but not the size of long-lived nanopores in the cell membrane. Biochim Biophys Acta **1848**: 958-966.

Pakhomov AG, Xiao S, Pakhomova ON, Semenov I, Kuipers MA, Ibey BL (2014) Disassembly of actin structures by nanosecond pulsed electric field is a downstream effect of cell swelling. Bioelectrochemistry **100**: 88-95.

Pakhomova ON, Gregory BW, Semenov I, Pakhomov AG (2013) Two modes of cell death caused by exposure to nanosecond pulsed electric field. PLoS One 8: e70278. DOI: 10.1371/journal.pone.0070278.

Pakhomova ON, Khorokhorina VÅ, Bowman AM, Rodaitė-Riševičienė R, Saulis G, Xiao S, Pakhomov AG (2012) Oxidative effects of nanosecond pulsed electric field exposure in cells and cell-free media. Arch Biochem Biophys **527**: 55-64.

Perdisa F, Gostyńska N, Roffi A, Filardo G, Marcacci M, Kon E (2015) Adipose-derived mesenchymal stem cells for the treatment of articular cartilage: a systematic review on preclinical and clinical evidence. Stem Cells Int **2015**: 597652. DOI: 10.1155/2015/597652

Polidori P, Lee S, Kauffman RG, Marsh BB (1999) Low voltage electrical stimulation of lamb carcasses: effects on meat quality. Meat Science **53**: 179-182.

Qin S, Yin H, Yang C, Dou Y, Liu Z, Zhang P, Yu H, Huang Y, Feng J, Hao J, Hao J, Deng L, Yan X, Dong X, Zhao Z, Jiang T, Wang HW, Luo SJ, Xie C (2016) A magnetic protein biocompass. Nat Mater **15**: 217-226.

Rems L, Miklavčič D, Pucihar G (2014) Induced transmembrane voltage during cell electrofusion using nanosecond electric pulses. In: Roa Romero L. (eds) XIII Mediterranean Conference on Medical and Biological Engineering and Computing 2013. IFMBE Proceedings, **41**. Springer, Cham, pp: 896-899.

Rems L, Ušaj M, Kandušer M, Reberšek M, Miklavčič D, Pucihar G (2013) Cell electrofusion using nanosecond electric pulses. Sci Rep **3**: 3382. DOI: 10.1038/srep03382.

Rossini P, Burke D, Chen R, Cohen L, Daskalakis Z, Di Iorio R, Di Lazzaro V, Ferreri F, Fitzgerald P, George M (2015) Non-invasive electrical and magnetic stimulation of the brain, spinal cord, roots and peripheral nerves: basic principles and procedures for routine clinical and research application. An updated report from an I.F.C.N. Committee. Clin Neurophysiol **126**: 1071-1107.

Rouabhia M, Park H, Meng S, Derbali H, Zhang Z (2013) Electrical stimulation promotes wound healing by enhancing dermal fibroblast activity and promoting myofibroblast transdifferentiation. PLoS One 8: e71660. DOI: 10.1371/journal.pone.0071660.

Schoellhammer CM, Blankschtein D, Langer R (2014) Skin permeabilization for transdermal drug delivery: recent advances and future prospects. Expert Opin Drug Deliv **11**: 393-407.

Schoenbach KH (2018) From the basic science of biological effects of ultrashort electrical pulses to medical therapies. Bioelectromagnetics **39**: 257-276.



Schoenbach KH, Hargrave SJ, Joshi RP, Kolb JF, Nuccitelli R, Osgood C, Pakhomov A, Stacey M, Swanson RJ, White JA (2007) Bioelectric effects of intense nanosecond pulses. IEEE **14**: 1088-1109.

Schoenbach KH, Joshi RP, Kolb JF, Chen N, Stacey M, Blackmore PF, Buescher ES, Beebe SJ (2004) Ultrashort electrical pulses open a new gateway into biological cells. IEEE **92**: 1122-1137.

Schwan HP (1956) Electrical properties of tissue and cell suspensions. Adv Biol Med Phys **5**: 147-209.

Semenov I, Xiao S, Pakhomov AG (2013) Primary pathways of intracellular Ca2+ mobilization by nanosecond pulsed electric field. Biochim Biophys Acta **1828**: 981-989.

Soba A, Suárez C, González MM, Cabrales LEB, Pupo AEB, Reyes JB, Tassé JPM (2018) Integrated analysis of the potential, electric field, temperature, pH and tissue damage generated by different electrode arrays in a tumor under electrochemical treatment. Math Comput Simul **146**: 160-176.

Sobrino-López A, Martín-Belloso O (2010) Review: potential of high-intensity pulsed electric field technology for milk processing. Food Eng Rev **2**: 17-27.

Somoza RA, Welter JF, Correa D, Caplan AI (2014) Chondrogenic differentiation of mesenchymal stem cells: challenges and unfulfilled expectations. Tissue Eng Part B Rev **20**: 596-608.

Tang X, Fan L, Pei M, Zeng L, Ge Z (2015) Evolving concepts of chondrogenic differentiation: history, state-of-the-art and future perspectives. Eur Cell Mater **30**: 12-27.

Teissie J, Tsong TY (1980) Evidence of voltageinduced channel opening in Na-K ATPase of humanerythrocyte membrane. J Membr Biol **55**: 133-140.

Toepfl S, Mathys A, Heinz V, Knorr D (2006) Potential of high hydrostatic pressure and pulsed electric fields for energy efficient and environmentally friendly food processing. Food Reviews International **22**: 405-423.

Tolstykh G, Beier H, Thompson G, Roth C, Ibey B (2013) Nanosecond pulsed electric fields activate intracellular signaling pathways. SPIE Newsroom. DOI: 10.1117/2.1201302.004736.

Ud-Din S, Bayat A (2014) Electrical stimulation and cutaneous wound healing: a review of clinical evidence. Healthcare (Basel) **2**: 445-467.

Ud-Din S, Sebastian A, Giddings P, Colthurst J, Whiteside S, Morris J, Nuccitelli R, Pullar C, Baguneid M, Bayat A (2015) Angiogenesis is induced and wound size is reduced by electrical stimulation in an acute wound healing model in human skin. PLoS One **10**: e0124502. DOI: 10.1371/journal.pone.0124502.

Uzieliene I, Bernotas P, Mobasheri A, Bernotiene E (2018) The role of physical stimuli on calcium channels in chondrogenic differentiation of mesenchymal stem cells. Int J Mol Sci **19**: E2998. DOI: 10.3390/ ijms19102998.

Vasilkoski Z, Esser AT, Gowrishankar T, Weaver JC (2006) Membrane electroporation: the absolute rate

equation and nanosecond time scale pore creation. Phys Rew E **74**: 021904.

Vernier PT, Sun YH, Marcu L, Salemi S, Craft CM, Gundersen MA (2003) Calcium bursts induced by nanosecond electric pulses. Biochem Biophys Res Commun **310**: 286-295.

Voldman J (2006) Electrical forces for microscale cell manipulation. Annu Rev Biomed Eng 8: 425-454.

Wang W, Wang ZY, Zhang GH, Clark CC, Brighton CT (2004) Up-regulation of chondrocyte matrix genes and products by electric fields. Clin Orthop Relat Res **427 Suppl**: S163-173.

Wang X, Li Y, He X, Chen S, Zhang JZ (2014) Effect of strong electric field on the conformational integrity of insulin. J Phys Chem A **118**: 8942-8952.

Wang Y, Rouabhia M, Lavertu D, Zhang Z (2017) Pulsed electrical stimulation modulates fibroblasts' behaviour through the Smad signalling pathway. J Tissue Eng Regen Med **11**: 1110-1121.

Weaver JC, Smith KC, Esser AT, Son RS, Gowrishankar T (2012) A brief overview of electroporation pulse strength-duration space: a region where additional intracellular effects are expected. Bioelectrochemistry **87**: 236-243.

Weaver JC, Vaughan TE, Martin GT (1999) Biological effects due to weak electric and magnetic fields: the temperature variation threshold. Biophys J **76**: 3026-3030.

Weinberg SH (2013) High-frequency stimulation of excitable cells and networks. PLoS One **8**: e81402. DOI: 10.1371/journal.pone.0081402.

Xu J, Wang W, Clark CC, Brighton CT (2009) Signal transduction in electrically stimulated articular chondrocytes involves translocation of extracellular calcium through voltage-gated channels. Osteoarthritis Cartilage **17**: 397-405.

Yamada M, Tanemura K, Okada S, Iwanami A, Nakamura M, Mizuno H, Ozawa M, Ohyama-Goto R, Kitamura N, Kawano M (2007) Electrical stimulation modulates fate determination of differentiating embryonic stem cells. Stem Cells **25**: 562-570.

Yang E, Li J, Cho M, Xiao S (2018) Cell fragmentation and permeabilization by a 1 ns pulse driven triplepoint electrode. Biomed Res Int **2018**: 4072983. DOI: 10.1155/2018/4072983.

Yao C, Hu X, Mi Y, Li C, Sun C (2009) Window effect of pulsed electric field on biological cells. IEEE **16**: 1259-1266.

Yarmush ML, Golberg A, Sersa G, Kotnik T, Miklavcic D (2014) Electroporation-based technologies for medicine: principles, applications, and challenges. Annu Rev Biomed Eng **16**: 295-320.

Yogesh K (2016) Pulsed electric field processing of egg products: a review. J Food Sci Technol **53**: 934-945.

Yu X, McGraw PA, House FS, Crowe JE, Jr. (2008) An optimized electrofusion-based protocol for generating virus-specific human monoclonal antibodies. J Immunol Methods **336**: 142-151.

Yuan X, Arkonac DE, Chao P-hG, Vunjak-Novakovic G (2014) Electrical stimulation enhances



cell migration and integrative repair in the meniscus. Sci Rep **4**: 3674. DOI: 10.1038/srep03674

Zeng C, Yang T, Deng Z-h, Yang Y, Zhang Y, Lei G-h (2015) Electrical stimulation for pain relief in knee osteoarthritis: systematic review and network meta-analysis. Osteoarthritis Cartilage **23**: 189-202.

Zhang K, Guo J, Ge Z, Zhang J (2014) Nanosecond pulsed electric fields (nsPEFs) regulate phenotypes of chondrocytes through Wnt/ β -catenin signaling pathway. Sci Rep **4**: 5836. DOI: 10.1038/srep05836.

Zhang X, Ren W, DeCaen P, Yan C, Tao X, Tang L, Wang J, Hasegawa K, Kumasaka T, He J (2012) Crystal structure of an orthologue of the NaChBac voltage-gated sodium channel. Nature **486**: 130-134.

Zhao M, Song B, Pu J, Wada T, Reid B, Tai G, Wang F, Guo A, Walczysko P, Gu Y, Sasaki T, Suzuki A, Forrester JV, Bourne HR, Devreotes PN, McCaig CD, Penninger JM (2006) Electrical signals control wound healing through phosphatidylinositol-3-OH kinase-gamma and PTEN. Nature **442**: 457-460.

Zimmermann U (1986) Electrical breakdown, electropermeabilization and electrofusion. Rev Physiol Biochem Pharmacol **105**: 175-256.

Editor's note: All comments/questions by the reviewers were answered by making changes in the text. Hence, there is no Discussion with Reviewers section.

The Scientific Editor responsible for this paper was Christine Hartmann.

